

Tasmannia insipida R.Br. Ex DC. Leaf Extracts Inhibit the Growth of the Acne Vulgaris Causing Bacterium *Cutibacterium acnes*

Ian Edwin Cock^{1,2,*}

¹School of Environment and Science, Nathan Campus, Griffith University, Brisbane, AUSTRALIA

²Centre for Planetary Health and Food Security, Nathan Campus, Griffith University, Brisbane, AUSTRALIA.

ABSTRACT

Introduction: Acne vulgaris is a skin conditions that affects most adolescents and also afflicts some adults. Medicinal plant extracts may provide leads for the development of new topical and/or oral therapies for acnes vulgaris, yet many traditional medicine plants are yet to be screened for growth inhibitory activity against *Cutibacterium acnes* (the major bacterial cause of acne). **Materials and Methods:** Methanolic and aqueous *Tasmannia insipida* R.Br. ex DC. Leaf extracts were investigated by disc diffusion and liquid dilution MIC assays against *Cutibacterium acnes*. Toxicity was determined using *Artemia franciscana* nauplii bioassays. **Results:** Methanolic and aqueous *T. insipida* leaf extracts displayed noteworthy bacterial growth inhibitory activity against *C. acnes* growth. The methanolic *T. insipida* leaf extract had particularly good antibacterial effects against *C. acnes*, with an LD MIC value of 752 µg/mL. Slightly higher LD MIC values were noted for the aqueous *T. insipida* leaf extract against *C. acnes* (LD MIC=1250 µg/mL). The methanolic and aqueous *T. insipida* leaf extracts were nontoxic in the *Artemia franciscana* bioassay, with LC₅₀ values substantially >1000 µg/mL. **Conclusion:** The lack of toxicity of the methanolic and aqueous *T. insipida* leaf extracts and their noteworthy inhibition of *C. acnes* growth indicate their potential to alleviate acne vulgaris. Further studies are warranted to isolate and identify the active components and to determine their antibacterial mechanism.

Keywords: Winteraceae, Brush Pepperberry, Australian Medicinal Plants, Acne Vulgaris, Skin Infection, Skin Inflammation, *Cutibacterium acnes*, *Artemia* lethality assay.

Correspondence:

Dr. Ian Edwin Cock

¹School of Environment and Science, and Centre for Planetary Health and Food Security, Nathan Campus, Griffith University, Brisbane, AUSTRALIA.

²Centre for Planetary Health and Food Security, Nathan Campus, Griffith University, Brisbane, AUSTRALIA.

Email: i.cock@griffith.edu.au

Received: 24-06-2025;

Revised: 07-08-2025;

Accepted: 13-10-2025.

INTRODUCTION

Acne vulgaris (generally referred to as acne) is an inflammatory skin condition, which is characterised by the presence of red scaly inflamed skin, pimples/pustules, painful subcutaneous lumps, whiteheads, blackheads and cystic lesions.¹ Acne is one on the most common infectious diseases globally. Indeed, it has been estimated that approximately 85% of the world's population are affected by acne vulgaris throughout their lives. Acne vulgaris can affect individuals of all ages although the incidence in 12-24 years olds is particular high.² Furthermore, individuals in that age group may suffer from longer and more severe outbreaks. Notably, prolonged acne can lead to anxiety and low self-esteem, which may result in social isolation and mental health issues. The incidence of acne vulgaris is influenced by a variety of factors, some of which can be controlled. Genetic factors (which cannot

be controlled) appear to be the most important indicator of acne vulgaris predisposition (particularly genetic predisposition for the production of high levels of androgens), accounting for ~80% of cases.² However, other controllable factors including diet, stress and cigarette smoking also contribute to the development of chronic acne vulgaris. Additionally, subdermal *Cutibacterium acnes* (previously called *Propionobacterium acnes*) infection also triggers acne vulgaris,³ providing a target to develop drugs for inhibiting the induction of acne vulgaris, as well as decreasing its severity.

Treatments for acne vary, depending on the severity of the condition. For mild cases, topical application of anti-acne creams are most frequently used, whereas both oral therapies and topical treatments may be used for severe cases of acne. The currently used medications include alpha hydroxyl acids, azelaic acid, benzoyl peroxide, isotretinoids, keratolytic soaps, retinoids and salicylic acid.⁴ Hormonal therapy (anti-androgens and/or androgen receptor antagonists) may also be used to treat severe acne. Additionally, antibiotics are useful in controlling chronic acne as they inhibit *C. acnes* infections, thereby blocking the development of inflammatory acne vulgaris symptoms.⁴



DOI: 10.5530/pc.2025.4.23

Copyright Information :

Copyright Author (s) 2025 Distributed under
Creative Commons CC-BY 4.0

Publishing Partner : Manuscript Technomedia, [www.mstechnomedia.com]

However, all of these remedies are associated with toxicity and unwanted side-effects. Additionally, prolonged use of broad-spectrum antibiotics is not recommended as it can result in the development of antibiotic-resistant *C. acnes* (as well as other pathogenic bacteria), especially to the commonly used tetracycline and macrolide classes of antibiotics.⁵ The use of complementary and alternative therapies that inhibit the growth of *C. acnes* is also relatively common in many regions of the world.^{6,7} The development of new acne vulgaris therapies from traditional plant-based medicines has potential and several studies have begun to screen traditional medicines against *C. acne*.⁷⁻⁹ However, many other plants are yet to be screened against *C. acnes* for growth inhibitory activity.

Tasmannia insipida R.Br. ex DC. (Family Winteraceae; commonly known as brush pepperberry) is a native Australian shrub that grows in cool wet coastal forests of eastern Australia. The First Australians used *T. insipida* leaves and berries as flavouring agents,^{10,11} although we were unable to find reports of the use of *T. insipida* for therapeutic uses. The related species *Tasmannia lanceolata* (Poir.) A.C.Sm. (commonly known as mountain pepperberry or Tasmanian pepperberry) was used as a medicine to treat a variety of conditions, including many illnesses caused by pathogenic bacteria.^{12,13} The antimicrobial properties of *T. insipida* have not been extensively studied and we only found a couple of studies that screened *T. insipida* leaf extracts against a limited number of bacterial pathogens.^{10,11} Notably, those studies also screened *Tasmannia stipitata* (Vickery) A.C.Sm. (Commonly known as Dorriggo pepper or northern pepperberry) and reported noteworthy antibacterial activity against several bacterial pathogens. A further study also reported good antibacterial activity for *T. stipitata* against some food spoilage bacteria.¹⁴ In contrast, the antibacterial properties of *T. lanceolata* leaves and berries have been substantially more extensively studied and noteworthy activity has been reported against a more extensive panel of bacteria.^{10,11,13} This study screened *T. insipida* solvent extractions for the ability to inhibit *C. acnes* growth, and quantified the antibacterial potency by MIC determination.

MATERIALS AND METHODS

Plant source and extraction

The *Tasmannia insipida* R.Br. ex DC. Leaves used in this study were provided by the Qld. Bushfoods Association, Australia as pre-dried and coarse milled whole plant material. The leaf material was stored at -30°C until use. The dried leaves were freshly ground to a coarse powder prior to extraction. Individual 1 g quantities of the ground fruit were weighed separately into tubes and 50 mL of methanol (AR grade, Ajax Fine Chemicals, Australia) or sterile deionised water were added. The fruits were extracted in each solvent for 24 hr at 4°C with gentle shaking. The extracts were filtered through filter paper (Whatman No. 54) under vacuum, followed by drying in a vacuum oven at 50°C. The

resultant dry extracts were weighed and redissolved in 10 mL of deionised water (containing 0.5% DMSO), and passed through a 0.22 µm filter (Sarstedt) to remove particulates. The extracts were then stored at 4°C until use.

Qualitative phytochemical studies

Phytochemical analysis of the *T. insipida* leaf extracts for the presence of cardiac glycosides, alkaloids, saponins, tannins, flavonoids, phenolic compounds, phytosterols, flavonoids and triterpenoids was achieved as previously described.¹⁵⁻¹⁷

Antibacterial screening

Test bacterial strains

All media and other materials was supplied by Oxoid Ltd., Australia unless otherwise specified. A reference strain of *Cutibacterium acnes* (ATCC6919) were purchased from American Type Culture Collection, USA. The *C. acnes* stock was cultured using a thioglycollate liquid media (Oxoid Ltd., Australia) under induced anaerobic conditions through the use of anaerobic jars and AnaeroGen™ 3.5 L atmospheric generation systems (Thermo Scientific) at 37°C for 72 hr.

Evaluation of antibacterial activity

Antibacterial activity screening of the *T. insipida* leaf extracts was achieved using modified disc diffusion assays.¹⁸⁻²⁰ Briefly, 100 µL of each individual bacteria was grown separately in 20 mL of broth until an approximate count of 10⁸ cells/mL was reached. A volume of 100 µL of the bacterial suspension was spread onto nutrient agar plates and the extracts were tested for antibacterial activity using 6 mm sterilised filter paper discs. Discs were infused with 10 µL of the individual extracts, allowed to dry, and placed onto the inoculated plates. The plates were left to stand at 4°C for 2 hr before incubation. The plates were incubated under induced anaerobic conditions at 37°C for 72 hr. The diameters of the inhibition zones (ZOIs) were measured to the closest whole millimetre. The assays were completed three times, each with internal triplicates (*n*=9). Mean values (±SEM) are reported in this study. Ampicillin (10 µg) and vancomycin (5 µg) discs were obtained from Oxoid Ltd., Australia and used as positive controls to compare antibacterial activity. Filter discs infused with 10 µL of distilled water (containing 0.5% DMSO) were used as negative controls.

Minimum inhibitory concentration (MIC) determination

The minimum inhibitory concentration for each extract was determined using two methods. A liquid dilution MIC assay was employed as it is generally considered the most sensitive bacterial growth inhibitory assay.²¹ Furthermore, as microplate liquid dilution (LD) MIC assays are perhaps the most commonly used method of quantifying bacterial growth inhibition efficacy, use of this method allows for comparisons with other studies. A solid

phase agar disc diffusion (DD) assay was also used in this study as this bioassay was deemed to provide a closer representation of the environment and conditions relevant to solid-phase skin systems.

Microplate liquid dilution MIC assay

The MICs of the extracts were evaluated by standard methods.^{21,22} Briefly, overnight bacterial cultures were added dropwise to fresh liquid broth and the turbidity was visually adjusted to produce a McFarlands number 1 standard culture. This was subsequently diluted 1 in 50 with fresh broth, resulting in the MIC assay inoculum culture. A volume of 100 µL sterile broth was added to all wells of a 96 well plate. Test extracts or control antibiotics (100 µL) were then added to the top row of each plate and 1 in 2 serial dilutions were prepared in each column of wells by transferring 100 µL from the top well to the next well in each column, etc. A growth control (without extract) and a sterile control (without inoculum) were included on each plate. A volume of 100 µL of bacterial culture inoculum was added to all wells except the sterile control wells. The inoculated plates were incubated under induced anaerobic conditions at 37°C for 72 hr. p-Iodonitrotetrazolium violet (INT) was obtained from Sigma, Australia and dissolved in sterile deionised water to prepare a 0.2 mg/mL INT solution. A 40 µL volume of this solution was added into all wells and the plates were incubated for a further 6 hr at 30°C. Following incubation, the MIC was visually determined as the lowest dose at which colour development was inhibited.

Disc diffusion MIC assay

The minimum inhibitory concentrations (MIC) of the extracts was also evaluated by disc diffusion assay as previously described.^{23,24} Briefly, the *T. insipida* leaf extracts were diluted in deionised water (containing 0.5% DMSO) and tested across a range of concentrations. Discs were infused with 10 µL of the extract dilutions, allowed to dry and placed onto inoculated plates. The assay was achieved as outlined above and graphs of the zone of inhibition versus Ln concentration were plotted. Determination of MIC values were achieved using linear regression.

Toxicity screening

Reference toxin for toxicity screening

Potassium dichromate ($K_2Cr_2O_7$) (AR grade, Chem-Supply, Australia) was prepared in deionised water (4 mg/mL) and serially diluted in artificial seawater for use in the *Artemia franciscana* nauplii bioassay.

Artemia franciscana nauplii toxicity screening

Toxicity was assessed using a modified *Artemia franciscana* nauplii lethality assay.^{25,26} Briefly, 400 µL of seawater containing 38 (mean 38.4, $n=125$, SD 13.7) *A. franciscana* nauplii were added to wells of a 48 well plate and immediately used in the bioassay.

A volume of 400 µL of the reference toxin or the diluted plant extracts were transferred to the wells and incubated at $25\pm1^\circ\text{C}$ under artificial light (1000 Lux). For each plate, a 400 µL seawater negative control was run in triplicate. The wells were assessed at regular intervals and the number of dead counted. The nauplii were deemed dead if no movement of the appendages was observed within 10 sec. After 24 hr, all nauplii were sacrificed and counted to determine the total % mortality per well. The LC_{50} with 95% confidence limits for each treatment was calculated using probit analysis.

Statistical analysis

Data is expressed as the Mean $\pm$ SEM of at least three independent experiments, each with internal triplicates ($n=9$).

RESULTS

Liquid extraction yields and qualitative phytochemical screening

1 g masses of the dried *T. insipida* leaf powder were extracted separately with methanol and water, resulting in yields of 118 and 104 mg respectively (Table 1). The extracts were dried and were subsequently resuspended in 10 mL of deionised water (containing 0.5% DMSO), resulting in the concentrations presented in Table 1. Qualitative phytochemical studies showed that the methanolic and aqueous *T. insipida* leaf extracts had similar phytochemical profiles. Both extracts contained high levels of phenolic compounds and flavonoids, as well as moderate levels of saponins and tannins.

Inhibition of *Cutibacterium acnes* growth

To determine the ability of the *T. insipida* leaf extracts to inhibit the growth of *C. acnes*, 10 µL of each extract was screened using a disc diffusion assay. Bacterial growth was inhibited by both the methanolic and aqueous *T. insipida* leaf extracts (Figure 1). The inhibition of *C. acnes* growth by the methanolic extract was particularly noteworthy (as judged by zone of inhibition (ZOI), with inhibition zones of 9.6 mm, compared to 8.3 mm for the aqueous extract. Indeed, the growth inhibition by the aqueous extracts was comparable to that of the ampicillin control (10 µg), which gave 10.3 mm ZOIs. In contrast, the vancomycin (5 µg) control produced larger zones of inhibition (11.5 mm), indicating stronger antibacterial activity. However, it is noteworthy that the antibiotic controls used in this study consisted of relatively high doses of pure antibiotics. In contrast, the extracts are crude mixtures, which would contain many individual compounds. Thus, it is likely that the growth inhibitory activity of individual bioactive extract component(s) is/are particularly promising. Therefore, the methanolic and aqueous *T. insipida* leaf extracts have potential for use in treating acne and further study is warranted.

Quantification of minimum inhibitory concentration (MIC)

The relative level of antimicrobial activity was further evaluated by determining the MIC values of the *T. insipida* leaf extracts against *C. acnes* (Table 2). Notably, substantial differences were evident between the results obtained in the disc diffusion (DD) assay and the liquid dilution (LD) MIC screening assays, with substantially lower MIC values recorded for the liquid dilution assay (752 and 1250 µg/mL for the methanolic and aqueous extracts respectively), compared to the solid phase assay (2208 and 3426 µg/mL respectively). These findings may reflect the classes of molecules in the extracts. Larger and/or lower polarity molecules do not readily diffuse through agar and therefore solid phase assays may provide erroneous results for these compounds. In contrast, larger and lower polarity molecules are more suitable for testing in liquid phase assays and these assays may provide a better understanding of their antibacterial activity. Given the substantially higher MIC values measured in the solid phase assays, it is likely that the bioactive compounds are relatively large and/or nonpolar, although this remains to be verified. Alternatively, potentiating compounds (with different physiochemical properties to the inhibitory compounds) may separate as they diffuse through the agar gel, whereas they will remain together in the liquid media assays, possibly accounting for these differences.

Quantification of toxicity

All extracts were screened in the *Artemia* nauplii assay at 2000 µg/mL (Figure 2). Additionally, potassium dichromate was also tested in the bioassay as a reference toxin. Potassium dichromate was rapid in its onset of mortality, promoting nauplii death within the first 3h of exposure, with 100% mortality evident within 5 hr (unpublished results). In contrast, the methanolic and aqueous *T. insipida* leaf extracts induced substantially <50% mortality following 24 hr exposure. As 24 hr LC₅₀ values >1000 µg/mL have previously been defined as nontoxic in this assay,^{25,26} the methanolic and aqueous *T. insipida* leaf extracts were deemed to be nontoxic and their LC₅₀ values were not further determined.

DISCUSSION

Acne vulgaris is one of the most common infectious diseases globally and the most common cause of chronic dermal inflammation globally, affecting approximately 85% of the population throughout their lives.² It is particularly common amongst people 12-24 years old and is often neglected by medical researchers as it has a low mortality rate. However, chronic acne vulgaris can lead to psychological issues, social isolation and emotional impairment amongst adolescents. Additionally, ineffective therapy may cause permanent scarring and may result in lifelong self-esteem issues and depression. Whilst there are multiple therapeutic options to control acne vulgaris, none are completely effective, and all are associated with toxicities and/or

Table 1: The mass of dried extracted material, the concentration after resuspension in deionised water (0.5 % DMSO) and qualitative phytochemical screenings of the *T. insipida* leaf extracts.

		Methanolic extract	Aqueous extract
Mass of extracted material (mg)		118	104
Concentration of resuspended extract (mg/mL)		12	10
Phenols	Total phenols	+++	+++
	Water soluble phenols	+++	+++
	Insoluble phenols	+++	+++
Saponins	Froth persistence	++	++
	Emulsion test	+	++
Cardiac glycosides	Keller-Kiliani Test	-	-
Triterpenoids	Salkowski Test	+	-
Phytosterols	Acetic Anhydride Test	+	-
Alkaloids	Meyer's Test	-	-
	Wagner's Test	-	-
	Draggendorff's Test	-	-
Flavonoids	Kumar Test	+++	+++
Tannins	Ferric Chloride Test	++	+++
	Lead Acetate Test	+	++
Anthraquinones	Free	-	-
	Combined	-	-

+++ indicates a large response; ++ indicates a moderate response; + indicates a minor response; - indicates no response in the assay.




Figure 1: Growth inhibitory activity of the *T. insipida* leaf extracts against *C. acnes* (ATCC 6919) measured as zones of inhibition (mm). M=methanolic extract; W=aqueous extract; Amp=ampicillin (10 µg); Van=vancomycin (5 µg); NC=negative control (0.5 % DMSO). Results are expressed as mean zones of inhibition±SEM. * indicates results that are significantly different to the negative control ($p < 0.05$).




Figure 2: The lethality of the *T. insipida* leaf extracts (2000 µg/mL), as well as the potassium dichromate (1000 µg/mL) and seawater controls towards *Artemia franciscana* nauplii after 24 hr exposure. M=methanolic extract; W=aqueous extract; PC=potassium dichromate control; NC=negative (seawater) control. * indicates results that are significantly different to the negative control ($p < 0.05$). Results are expressed as mean % mortality±SEM.

Table 2: Disc diffusion and liquid dilution MICs against *P. acnes* growth (µg/mL) of the *T. insipida* extracts.

Extract	MIC (µg/mL)	
	DD	LD
Methanolic extract	2208	752
Aqueous extract	3426	1250

DD disc diffusion; LD liquid dilution; Numbers indicate the mean DD MIC and LD MIC values of triplicate determinations. - indicates no inhibition at any concentration tested.

unwanted side-effects.^{3,4} There is a need to develop safer and more effective therapies that can be used for topical and oral treatment to inhibit the growth of *C. acnes*. Traditional plant-based medicines are especially promising and an examination of ethnobotanical records can highlight promising plants (and plant combinations) for screening. Many traditional medicine plant have been used for hundreds (or even thousands) of years to inhibit bacterial growth, and in some case their efficacy has been verified by rigorous scientific examination. Additionally, the use of natural alternatives to inhibit the growth of acne vulgaris-inducing bacteria may be more acceptable to consumers due to their natural origin and consumer perception of safety.

Multiple plant species have been used traditionally to treat acne, including (but not limited to) *Allium cepa* L. (onion), *Allium sativum* L. (garlic), *Aloe vera* L., *Camellia senensis* (L.) Kuntze (tea plant), *Cannabis sativa* L. (cannabis, marijuana), *Echinacea purpurea* (echinacea), *Eucalyptus globus* Labill. (Blue gum), *Lavendula angustifolia* Mill. (Lavender), *Portulaca oleraceae* L. (common purslane), *Salvia officinalis* Spenn. (Rosemary), and *Thyme vulgaris* L. (thyme).²⁷ Several of these have already been tested for the ability to inhibit the growth of *C. acnes*, although the growth inhibitory properties of many of these are yet to be

verified. Substantially fewer studies have screened Australian plants for the ability to inhibit the growth of *C. acnes*, although several recent studies have begun to examine the potential of Australian plants to inhibit *C. acnes* growth. In particular, *Terminalia ferdinandiana* Exell. Fruit extracts were recently reported to be good inhibitors of *C. acnes* growth (as well as several other skin pathogens).^{28,29} Similarly, the anti-*C. Acnes* activity of several Australian *Syzygium* spp.³⁰ and *Acronychia acidula* F. Muell. Fruit extracts has also been documented.³¹

This study examined the growth inhibitory properties of methanolic and aqueous *T. insipida* leaf extracts against *C. acne* (the main bacterial cause of acne vulgaris). *Tasmannia insipida* leaf extracts were selected for screening in this study due to the taxonomic relationship of this species with *T. stipitata* and *T. lanceolata*, which have both previously been reported to have noteworthy antibacterial activity.¹⁰⁻¹³ Our study confirmed the potential of the *T. insipida* leaf extracts for inhibiting the growth of the main acne vulgaris causing bacteria, *C. acnes*. The methanolic extract was the most promising growth inhibitor, with an LD MIC value of 752 µg/mL measured, compared to an LD MIC of 1250 µg/mL for the aqueous extract. Studies using extracts prepared from other plants have reported comparable MIC values as signifying potent inhibitory activity. Extracts produced from *T. ferdinandiana* leaves were reported to be potent inhibitors of *C. acnes* growth, with an MIC value of 625 µg/mL.²⁸ Therefore, the *T. ferdinandiana* leaf extracts are only slightly stronger than the methanolic *T. insipida* extract tested in our study. Another study screened methanolic and aqueous *Acronychia acidula* F. Muell. (Commonly known as lemon aspen) against *C. acnes* and reported noteworthy inhibitory activity for the aqueous extract (MIC=1455 µg/mL), although the methanolic extract was completely ineffective against that bacterium.³¹ Notably, the

substantially lower MIC value of the methanolic *T. insipida* leaf extract screened in our study indicates that it is more potent than the *A. acidula* extracts tested in the earlier study. Thus, the use of the methanolic *T. insipida* leaf extract to treat acne vulgaris is promising and further testing is warranted.

Whilst the antibacterial components of the *T. insipida* leaf extracts were not identified in this study, a several classes of compounds were highlighted by the qualitative phytochemical analysis studies. The detection of moderate to high levels of polyphenolics, tannins and flavonoids was particularly noteworthy. Many studies have reported potent antibacterial activities for a wide variety of flavonoids.³² This has been attributed to a variety of mechanisms, including their ability to complex with extracellular and soluble proteins, as well as bacterial cell wall proteins.³³ Similarly, multiple tannins have broad-spectrum antibacterial activity via a variety of intra and extracellular mechanisms, including the precipitation of microbial proteins.³⁴ Polyphenolics are toxic to microorganisms via enzyme inhibition mechanisms, possibly through non-specific interaction with proteins, or by reaction with sulphhydryl groups.³⁵ It is also likely that other phytochemical classes may also contribute to the growth inhibitory properties of these extracts. Further studies to elucidate the phytochemicals in this extract (and their potential therapeutic mechanisms) are required.

Notably, the methanolic and aqueous *T. insipida* leaf extracts tested in this study were both nontoxic towards *Artemia* nauplii and are therefore likely to be safe for topical application, as well for oral usage. However, further toxicity studies using human cell lines (and subsequent *in vivo* studies) are required to confirm the safety of these extracts before they are accepted as natural therapies for acne vulgaris. Furthermore, whilst our study reported the *T. insipida* leaf extracts to be nontoxic, it is noteworthy that these studies have only examined acute toxicity. Pharmacodynamic and pharmacokinetic studies are required to determine the ability of the extract components to cross the skin barrier, their duration in the blood stream prior to clearance, and the urinary excretory products. Indeed, such studies are required for any formulation to ensure that their components do not accumulate and cause chronic toxicity.

CONCLUSION

The results of this study demonstrate the potential of the *T. insipida* leaf extracts as natural antibacterial components to treat acne vulgaris. The methanolic extract was particularly potent and has particular promise for acne vulgaris drug discovery. Furthermore, the lack of toxicity of the extract indicates the suitability of that extract for topical or oral use.

ACKNOWLEDGEMENT

Financial support for this work was provided by Centre for Planetary Health and Food Security, Griffith University.

ABBREVIATIONS

DMSO: Dimethyl sulfoxide; **LC₅₀:** The concentration required to achieve 50% mortality; **MIC:** Minimum inhibitory concentration; **ZOI:** Zone of inhibition.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

SUMMARY

- Methanolic and aqueous *T. insipida* leaf extracts were screened for the ability to block the growth of *Cutibacterium acnes*.
- The methanolic *T. insipida* leaf extract was a particularly good inhibitor of *C. acnes* growth (LD MIC=752 µg/mL).
- The aqueous *T. insipida* leaf extract was also a good inhibitor of *C. acnes* growth, albeit with a higher LD MIC value (LD MIC=1250 µg/mL).
- The nontoxicity of the *T. insipida* leaf extracts was verified using the *Artemia* nauplii toxicity bioassay.

REFERENCES

1. Mayo Clinic. Acne. <https://www.mayoclinic.org/diseases-conditions/acne/symptoms-causes/syc-20368047>. Cited 14 March 2022.
2. Lambrechts IA, de Canha MN, Lall N. Exploiting medicinal plants as possible treatments for acne vulgaris. In Medicinal Plants for Holistic Health and Well-Being, 2018; Academic Press, USA:117-43.
3. Dréno B. What is new in the pathophysiology of acne, an overview. Journal of the European Academy of Dermatology and Venereology 2017;31:8-12.
4. Nasri H, Bahmani M, Shahinfard N, et al. Medicinal plants for the treatment of acne vulgaris: a review of recent evidences. Jundishapur Journal of Microbiology 2015;8(11):e25580.
5. Cheesman MJ, Ilanko A, Blonk B, et al. Developing new antimicrobial therapies: are synergistic combinations of plant extracts/compounds with conventional antibiotics the solution?. Pharmacognosy Reviews 2017;11(22):57-72.
6. De Wet H, Nciki S, van Vuuren SF. Medicinal plants used for the treatment of various skin disorders by a rural community in northern Maputland, South Africa. Journal of Ethnobiology and Ethnomedicine 2013;9(1):51.
7. Kumar GS, Jayaveera KN, Kumar CK, et al. Antimicrobial effects of Indian medicinal plants against acne-inducing bacteria. Tropical Journal of Pharmaceutical Research 2007;6(2):717-23.
8. Orchard A, Van Vuuren SF, Viljoen AM, et al. The *in vitro* antimicrobial evaluation of commercial essential oils and their combinations against acne. International Journal of Cosmetic Science 2018;40(3):226-43.
9. Ali-Shtayeh MS, Al-Assali AA, Jamous RM. Antimicrobial activity of Palestinian medicinal plants against acne-inducing bacteria. African Journal of Microbiology Research 2013;7(21):2560-73.
10. Cock IE, Winnett V, Sirdaarta J, et al. The potential of selected Australian medicinal plants with anti-*Proteus* activity for the treatment and prevention of rheumatoid arthritis. Pharmacognosy Magazine 2015; 11(42 Suppl 1):S190-S208.
11. Winnett V, Sirdaarta J, White A, et al. Inhibition of *Klebsiella pneumoniae* growth by selected Australian plants: Natural approaches for the prevention and management of ankylosing spondylitis. Inflammopharmacology 2017;25(2):223-35.
12. Cock IE. The phytochemistry and chemotherapeutic potential of *Tasmannia lanceolata* (Tasmanian pepper): A review. Pharmacognosy Communications 2013;3(4):13-25.
13. Winnett V, Boyer H, Sirdaarta J, Cock IE. The potential of *Tasmannia lanceolata* as a natural preservative and medicinal agent: Antimicrobial activity and toxicity. Pharmacognosy Communications 2014;4(1):42-52.
14. Hart C, Ilanko P, Sirdaarta J, et al. *Tasmannia stipitata* as a functional food/natural preservative: Antimicrobial activity and toxicity. Pharmacognosy Communications 2014;4(4):33-47.

15. Courtney R, Sirdaarta J, Matthews B, *et al.* Tannin components and inhibitory activity of Kakadu plum leaf extracts against microbial triggers of autoimmune inflammatory diseases. *Pharmacognosy Journal* 2015;7(1):18-31. DOI: 10.5530/pj.2015.7.2
16. Murhekar S, Wright MH, Greene AC, *et al.* Inhibition of *Shewanella* spp. growth by *Syzygium australe* and *Syzygium leuhmannii* extracts: Natural methods for the prevention of fish spoilage. *Journal of Food Science and Technology*;54(10):3314-26.
17. Ilanko A, Cock IE. The interactive antimicrobial activity of conventional antibiotics and *Petalostigma* spp. extracts against bacterial triggers of some autoimmune inflammatory diseases. *Pharmacognosy Journal* 2019;11(2):292-309.
18. Hutchings A, Cock IE. An interactive antimicrobial activity of *Embelica officinalis* Gaertn. Fruit extracts and conventional antibiotics against some bacterial triggers of autoimmune inflammatory diseases. *Pharmacognosy Journal* 2018;10(4):654-62.
19. Ilanko P, McDonnell PA, van Vuuren S, *et al.* Interactive antibacterial profile of *Moringa oleifera* Lam. extracts and conventional antibiotics against bacterial triggers of some autoimmune inflammatory diseases. *South African Journal of Botany* 2019;124:420-35.
20. Cock IE. Antimicrobial activity of *Syzygium australe* and *Syzygium leuhmannii* leaf methanolic extracts. *Pharmacognosy Communications* 2012;2(2):71-7.
21. Hübsch Z, Van Zyl RL, Cock IE, *et al.* Interactive antimicrobial and toxicity profiles of conventional antimicrobials with Southern African medicinal plants. *South African Journal of Botany* 2014;93:185-97.
22. Tiwana G, Cock IE, White A, *et al.* Use of specific combinations of the triphala plant component extracts to potentiate the inhibition of gastrointestinal bacterial growth. *Journal of Ethnopharmacology* 2020;260:112937. DOI: 10.1016/j.jep.2020.112937
23. Wright MH, Matthews B, Arnold MS, *et al.* The prevention of fish spoilage by high antioxidant Australian culinary plants: *Shewanella putrefaciens* growth inhibition. *International Journal of Food Science and Technology* 2016;51(3):801-13.
24. Cock IE. Antimicrobial activity of *Syzygium australe* and *Syzygium leuhmannii* leaf methanolic extracts. *Pharmacognosy Communications* 2012;2(2):71-7.
25. Cock IE, Ruebhart DR. Comparison of the brine shrimp nauplii bioassay and the ToxScreen-II test for the detection of toxicity associated with *Aloe vera* (*Aloe barbadensis* Miller) leaf extract. *Pharmacognosy Research* 2009;1(2):98-101.
26. Ruebhart DR, Wickramasinghe W, Cock IE. Protective efficacy of the antioxidants vitamin E and Trolox against *Microcystis aeruginosa* and microcystin-LR in *Artemia franciscana* nauplii. *Journal of Toxicology and Environmental Health, Part A* 2009;72(24):1567-75.
27. Tabassum N, Hamdani M. Plants used to treat skin diseases. *Pharmacognosy Reviews* 2014;8(15):52-60.
28. McManus K, Wood A, Wright MH, *et al.* *Terminalia ferdinandiana* Exell. extracts inhibit the growth of body odour forming bacteria. *International Journal of Cosmetic Science* 2017;39(5):500-10.
29. Cock IE, Wright MH, Matthews B, *et al.* Bioactive compounds sourced from *Terminalia* spp. in bacterial malodour prevention: an effective alternative to chemical additives. *International Journal of Cosmetic Science* 2019;41(5):496-508. DOI: 10.1111/ics.12567
30. Wood A, McManus KDB, Wright MH, *et al.* Growth inhibitory activity of selected Australian *Syzygium* species against malodour forming bacteria. *Pharmacognosy Communications* 2017;7(3):129-36.
31. Wright MH, Wood A, Greene T, *et al.* Growth inhibitory activity of *Acronychia acidula* F. Muell. fruit extracts towards malodour-forming bacteria. *Pharmacognosy Communications* 2020;10(2):95-101
32. Narayana KR, Reddy MS, Chaluvadi MR, *et al.* Bioflavonoids classification, pharmacological, biochemical effects and therapeutic potential. *Indian Journal of Pharmacology* 2001;33(1):2-16.
33. Kaur GJ, Arora DS. Antibacterial and phytochemical screening of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi*. *BMC Complementary and Alternative Medicine* 2009;9:1.
34. Buzzini P, Arapitsas P, Goretti M, *et al.* Antimicrobial activity of hydrolysable tannins. *Medicinal Chemistry* 2008;8:1179-87.
35. Cowan MM. Plant products as antimicrobial agents. *Clinical Microbiology Reviews* 1999;12:564-82.

Cite this article: Cock IE. *Tasmannia insipida* R.Br. Ex DC. Leaf Extracts Inhibit the Growth of the Acne Vulgaris Causing Bacterium *Cutibacterium acnes*. *Pharmacognosy Communications*. 2025;15(4):179-85.