

# Research Article

## Renal and Hepatic Antioxidant Status of Hyperglycemic Rats Treated with Single and Combinatorial Herbal Formulations.

Okey Alphosus Ojiako<sup>1</sup>, Paul Chidoka Chikezie<sup>2</sup>, Agomuo Chizaramoku Ogbuji<sup>3</sup>

<sup>1</sup>Department of Biochemistry, Federal University of Technology, Owerri, Nigeria

<sup>2</sup>Department of Biochemistry, Imo State University, Owerri, Nigeria

<sup>3</sup>Department of Food Science and Technology, Abia State Polytechnic, Aba, Nigeria

### ABSTRACT

**Background/Aim:** The present study ascertained the comparative capacities of single and combinatorial herbal formulations of *Acanthus montanus*, *Emilia coccinea*, *Hibiscus rosasinensis* and *Asystasia gangetica* to exert glycemic control and ameliorate oxidative stress in renal and hepatic tissues of alloxan-induced hyperglycemic rats (HyGR) following 30 days treatment. **Materials and Methods:** Phytochemical composition of the herbal extracts and fasting plasma glucose concentration (FPGC) of the rats were measured by standard methods. The post mitochondrial supernatant fractions (PMSF) of renal and hepatic tissues homogenates were measured for activity levels of glutathione peroxidase (GPOx), superoxide dismutase (SOD) and catalase (CAT) as well as glutathione (GSH) concentration. **Results:** Comparatively, flavonoids were the most abundant phytochemical present in the four herbal extracts. FPGC of treated HyGR was within the range of  $66.30 \pm 0.71$ - $133.00 \pm 0.63$  mg/dL compared to the untreated hyperglycemic rat group =  $368.20 \pm 0.72$  mg/dL ( $p < 0.05$ ). At the end of the experimental time, treated HyGR liver SOD activities were lower than that of normal group ( $p < 0.05$ ). Liver and kidney GPOx activities of HyGR were 2.36 folds and 3.93 folds lower than the normal rats. Furthermore, liver and kidney CAT activities of the treated HyGR were significantly ( $p < 0.05$ ) lower than the normal rats, whereas liver and kidney GSH/GSSG ratios were significantly ( $p < 0.05$ ) increased when compared with treated HyGR. **Conclusion:** Combinatorial herbal formulations exerted greater glycemic control than the single herbal formulations and caused a sparing effect on renal and hepatic antioxidants against reactive oxygen species.

**Key words:** Antioxidant, Hyperglycemia, Hepatic, Phytochemicals, Renal.

### INTRODUCTION

*Acanthus montanus* (Nees) T. Anderson belongs to the family of *Acanthaceae* and is widespread in Africa, the Balkans, Romania, Greece and Eastern Mediterranean. The plant is a thinly branched perennial shrub with basal clusters of oblong and lance shaped glossy, dark green leaves that measure about 30 cm long. *A. montanus* is commonly

known as mountain thistle or Devil's fig or Bear's Breech. The medicinal usefulness of the plant extracts have been previously reported elsewhere.<sup>1,2</sup> *Asystasia gangetica* L. T. Anderson is also known as creeping foxglove and belongs to the family of *Acanthaceae*. It is a fast-growing herbaceous weed of height between 0.3 – 0.4 m. Decoctions of *A. gangetica* are used in African and Asian ethno medicinal practices for the alleviation of stomach ache, urethral discharge, inflammation, rheumatism and dry cough. Also, *A. gangetica* is used for the management of diabetes mellitus (DM) and epilepsy as well as antidote to snake bite. *Emilia coccinea* (SIMS) G. Donis (family: *Aster- acca*) is commonly found in African countries like Congo,

\*Correspondence author:

**Dr. Paul Chidoka Chikezie**

Department of Biochemistry,

Imo State University, Owerri, Nigeria

E-mail: p\_chikezie@yahoo.com

DOI : 10.5530/pc.2015.2.7

Sudan, Kenya, Tanzanian, Malawi, Uganda and Nigeria.<sup>3</sup> In parts of Africa, this multipurpose plant serves as delicacies and herbal concoctions for remedies of numerous diseases including DM.<sup>3,4</sup> *Hibiscus rosasinensis* L. is an ever-green shrub that grows up to 1.8 – 2.4 m high. The family, *malvaceae*, is consist of 82 genera and over 1,500 species of which over 200 species are found in tropical and sub-tropical regions of the world. The flowers are hermaphrodites and vary in color depending on the species and location. The medicinal properties of *H. rosasinensis* have been reported by previous researchers.<sup>5-7</sup>

Hyperglycemia, which is one of the prominent defining features of DM, has been implicated in promoting overwhelming levels of reactive oxygen and nitrogen species (RONS)<sup>8,9</sup> and protein glycation.<sup>10,11</sup> Hyperglycemia mediated generation and accumulation of RONS may occur through one or combinations of the following molecular events: distortions of glycolytic pathways,<sup>12,13</sup> intercellular activation of sorbitol (polyol) pathways,<sup>14</sup> auto-oxidation of glucose<sup>15</sup> and non-enzymatic proteins and amino acids glycation.<sup>16,17</sup> Antioxidant systems prevent cellular damage from the adverse effects of xenobiotics, carcinogens and toxic RONS. Several reports showed that there are several biologically derived compounds with antioxidant functions.<sup>18-20</sup> Glutathione peroxidases (GPOx; EC 1.11.1.19) are ubiquitous selenium-containing peroxidases that catalyze the reduction of a variety of hydroperoxides (ROOH and H<sub>2</sub>O<sub>2</sub>) using GSH as co-factor. GPOx activity, among other functions, protects mammalian cells against oxidative damage. There are at least five GPOx isoenzymes found in mammals with the levels of each isoform vary depending on the tissue type.<sup>18,21</sup> Superoxide dismutase (SOD; EC 1.15.1.1) is metallo-enzyme that defines the basis for their classification; Mn-SOD<sup>22</sup> and Cu, Zn-SOD<sup>20,23</sup> Ni-SOD.<sup>24</sup> SOD activity neutralizes superoxide by transforming oxygen derived reactive species to peroxide, which can in turn be eliminated by the actions of catalase or GPOx activities. Catalase (CAT; EC 1.11.1.6) is a tetrameric haemin-enzyme consisting of four identical tetrahedrally arranged subunits of  $\approx$  60 kDa.<sup>18</sup> CAT activities prevents H<sub>2</sub>O<sub>2</sub>-induced cellular damage by converting H<sub>2</sub>O<sub>2</sub> to H<sub>2</sub>O and O<sub>2</sub>. To a large extent, the thiol-redox status of the cell is defined by cellular levels of GSH.<sup>25,26</sup> In addition, the reduced glutathione (GSH)/oxidized glutathione (GSSG) ratio (GSH:GSSG) is an important biomarker of redox environment in biologic systems.<sup>25,27</sup>

In modern medicine and traditional systems, medicinal plants have continued to provide valuable therapeutic agents for the treatment and management of diseases and disorders. Moreover, in spite of the higher acceptability

level of orthodox anti-diabetic therapeutics, the use several plants as herbal remedies by traditional healers for the treatment of diabetes mellitus is still very common especially in Africa and Asia continents.<sup>28-33</sup> Traditional herbs are commonly used in the form multi-herb recipe, which is believed to possess better pharmacological efficacy than the individual composite herb extracts. In that regard, the present study ascertained the comparative capacities of single and combinatorial herbal formulations of four medicinal plant extracts of *A. montanus*, *E. coccinea*, *H. rosasinensis* and *A. gangetica* to exert glycemic control in alloxan-induced hyperglycemic rats (HyGR). In addition, hyperglycemia causes glucose auto-oxidation mediated oxidative organ damage. Previous reports have revealed that cellular levels of the antioxidant system could serve as a reliable biomarker to establishing the pathogenesis and progression of DM especially with respect to tissue damage.<sup>34,35</sup> Therefore, studies were carried out to ascertain levels of renal and hepatic tissues derived primary oxidative stress indicators (GPOx, SOD, CAT activities and GSH/GSSG ratio) and their correlations, following the administration of the herbal formulations. The outcome of the present investigations will serve to give insights into the capacities of the various herbal formulations to ameliorate hyperglycemia and tissue derived oxidative stress in DM.

## MATERIALS AND METHODS

### Collection and preparation of herbal samples

Fresh leaves of *Acanthus montanus* (Nees) T. Anderson (ACMO), *Emilia coccinea* (SIMS) G. Don (EMCO) and *Hibiscus rosasinensis* L. (HIRO) were collected from uncultivated lands in Umuamacha Ayaba Umaeze, Osioma Ngwa LGA, Abia State, Nigeria. Fresh leaves of *Ayista-sia gangetica* L. T. Anderson (ASGA) were collected from Ubowuala, Emekuku, Owerri North LGA, Imo State, Nigeria. The four herbs were identified and authenticated by Dr. M. Ibe, School of Agriculture and Agricultural Technology (SAAT), Federal University of Technology, Owerri. All the leaves were collected between the months of July and August, 2009.

The leaves of individual plants were washed with a continuous flow of distilled water for 15 min and allowed to dry at laboratory ambient temperature (24  $\pm$  5°C). A 500 g part of each herbal samples were weighted using a triple beam balance (OHAU 750-50: Burlington, NC, USA) and dried in an oven (WTC BINDER, 7200 Tuttlingen, Germany) at 60°C until a constant weight was achieved. The dried leaves were packaged in dark polyethylene bags and kept in cold room (7  $\pm$  3°C) for 24 h before

pulverization. Next, the separate dried leaves were pulverized using Thomas-Willey milling machine (ASTM D-3182, INDIA), after which the ground samples were stored in air-tight plastic bottles with screw caps pending extraction.

### Extraction of herbal samples

Portions of 40 g of each pulverized dried samples of *A. montanus*, *E. coccinea*, *H. rosasinensis* and *A. gangetica* were subjected to repeated soxhlet extraction cycles for 2 h using 96% C<sub>2</sub>H<sub>5</sub>OH (BDH, U.K) as solvent to obtain a final volume of 500 mL of each herbal extract. These volumes of the extracts were concentrated and recovered in a rotary evaporator for 12 h at 60°C under reduced pressure. The extracts were dried in a desiccators for 24 h, wrapped in aluminum foil and stored in air-tight plastic bottles with screw caps at ≤4°C. The yields were calculated to be as follows: *A. montanus*=16.35% (*w/w*), *E. coccinea*=17.99% (*w/w*), *H. rosasinensis*=17.23% (*w/w*) and *A. gangetica* = 16.69% (*w/w*). The separate extracts were reconstituted in phosphate buffered saline (PBS) solution (extract vehicle), osmotically equivalent to 100 g/L PBS (90.0 g NaCl, 17.0 g Na<sub>2</sub>HPO<sub>4</sub>·2H<sub>2</sub>O and 2.43 g NaH<sub>2</sub>PO<sub>4</sub>·2H<sub>2</sub>O), before appropriated doses were administered to the experimental animals. Portions of the individual extracts were also measured for phytochemical contents.

### Phytochemical composition of herbal extracts

Flavonoids content was measured.<sup>36</sup> Five grams (5.0 g) of the herbal extract was weighted into 250 mL beaker, mixed with 100 mL of 96% C<sub>2</sub>H<sub>5</sub>OH for 30 min at 24±5°C and filtered using the Whatman No 1 filter paper. The filtrate was dried in an oven to constant weight and cooled in a dessicator.

The concentration of alkaloids of the herbal extracts was measured by the methods of Harborne.<sup>37</sup> A 5.0 g extract was weighted into 250 mL beaker containing 100 mL of 10% CH<sub>3</sub>COOH (Sigma, St. Louis, USA) in C<sub>2</sub>H<sub>5</sub>OH. The content was mixed thoroughly by vortex and allowed to stand with continuous agitation (Shaking Bath; Precision Scientific Inc., Chicago, IL) at 24±5°C for 4 h. The suspension was separated by simple filtration method using Whatman No 1 filter paper and concentrated by evaporation to ¼ of its original volume in a steam bath. Concentrated NH<sub>4</sub>OH (BDH, U.K) was subsequently added in droplets to the filtrate to form precipitates, which were subsequently separated by filtration. The precipitates were washed with 1% NH<sub>4</sub>OH, dried in an oven to constant weight and cooled in a dessicator.

Measurement of saponin content of the herbal extracts was by the methods of Harborne,<sup>37</sup> as reported by Oba-

doni and Ochuka.<sup>38</sup> Five grams (5.0 g) of the sample was mixed with 200 mL of 20% C<sub>2</sub>H<sub>5</sub>OH. The suspension was heated in a water bath at 55°C while stirring for 4 h and allowed to cool to 24 ± 5°C. The precipitate was harvested by simple filtration methods using Whatman No 42 filter paper. The filtrate was concentrated by reducing the total filtrate volume to 40 mL by evaporation in a steam bath. In a 250 mL capacity separating funnel, the 40 mL concentrated filtrate was mixed properly with 20 mL of C<sub>2</sub>H<sub>5</sub>OC<sub>2</sub>H<sub>5</sub> (BDH, U.K). The aqueous layer was recovered, whereas the organic layer was decanted. The purification and separation procedure was repeated twice. Finally, 60 mL of n-C<sub>4</sub>H<sub>9</sub>OH (BDH, U.K) was added to the extract followed by 10 mL of 5% NaCl (BDH, U.K) to precipitate saponins. The solution was evaporated to dryness in a water bath, dried in an oven and cooled in a dessicator.

### Calculations

The concentrations of flavonoids, alkaloids and saponin (mg/100 g sample weight) in the various herbal extracts were calculated thus:

$$C_{FAS} = \frac{(W2 - W1) \times 100}{Wt}$$

Where

C<sub>FAS</sub>: concentrations of flavonoids, alkaloids and saponins.

W2: weights of filter paper + precipitate.

W1: weight of filter paper.

Wt: weights of samples.

The Van-Burden and Robinson,<sup>39</sup> method as reported<sup>40</sup> was used to measure the concentration of tannins of the herbal extracts. A 5 g of the sample was mixed with 50 mL of distilled water for 1 h in a 50 mL capacity plastic bottle on a mechanical shaker. The mixture was filtered with Whatman No 42 filter paper. A 5 mL aliquot of the filtrate was mixed with 2 mL of 0.1 M FeCl<sub>3</sub> in 0.1 N HCl and 0.008 M K<sub>4</sub>(Fe)CN<sub>6</sub> (Sigma, St. Louis, USA) in a test tube. The absorbance of the analyte was measured at maximum absorptivity (λ<sub>max</sub>)=605 nm within 10 min and compared with the standards.

### Experimental animals

Male albino (Wistar) rats weighing between 150-160 g were maintained at room temperatures of 24±5°C, 30-55% of relative humidity on a 12-h light/12-h dark

cycle, with access to water and standard commercial feed (SCF) (Ewu Feed Mill, Edo State, Nigeria) *ad libitum* for 2 weeks acclimatization period. The handling of the animals was in accordance with the standard principles of laboratory animal care of the United States National Institutes of Health (NIH, 1978).

### Induction of diabetes/experimental design

Hypoglycemia was induced in the rats by single intra-peritoneal (i.p) injection of 90 mg/kg bw of alloxan monohydrate (Sigma, St. Louis, USA) in PBS solution (pH=7.4). The animals with fasting plasma glucose concentration (FPGC)>110 mg/dL for 5 consecutive days were considered diabetic and selected for the study. A total of 102 male Wistar rats were allotted into seventeen (17) groups of six (6) rats each. The animals were deprived of food and water for additional 16 h before commencement of treatment as described elsewhere.<sup>41</sup> The animal groups were designated on the basis of treatments received at regular intervals of 2 days for 30 days. Herbal treatments of the HyGR were defined as single herbal formulations (SHF): (HrACMO, HrASGA, HrEMCO and HrHIRO), double herbal formulations (DHF): (HrAGAM, HrAGEC, HrAGHR, HrAMEC, HrAMHR and HrECHR), triple herbal formulations (THF): (HrAGEH, HrAMAE, HrAMAH and HrAMEH) and quadruple herbal formulation (QHF): (HrAAEH).

- NORM: Normal rats received SCF + water *ad libitum* + 1.0 mL/kg of PBS.
- DIAB: HyGR received SCF + water *ad libitum* + 1.0 mL/kg of PBS.
- HrACMO: HyGR received SCF + water *ad libitum* + *A. montanus* (20 mg/kg in PBS; i.p.).
- HrASGA: HyGR received SCF + water *ad libitum* + *A. gangetica* (20 mg/kg in PBS; i.p.).
- HrEMCO: HyGR received SCF + water *ad libitum* + *E. coccinea* (20 mg/kg in PBS; i.p.).
- HrHIRO: HyGR received SCF + water *ad libitum* + *H. rosasinensis* (20 mg/kg in PBS; i.p.).
- HrAGAM: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1 *w/w*) of *A. gangetica* + *A. montanus* (20 mg/kg in PBS; i.p.).
- HrAGEC: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1 *w/w*) of *A. gangetica* + *E. coccinea* (20 mg/kg in PBS; i.p.).
- HrAGHR: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1 *w/w*) of *A. gangetica* + *H. rosasinensis* (20 mg/kg in PBS; i.p.).

- HrAMEC: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1 *w/w*) of *A. montanus* + *E. coccinea* (20 mg/kg in PBS; i.p.).
- HrAMHR: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1 *w/w*) of *A. montanus* + *H. rosasinensis* (20 mg/kg in PBS; i.p.).
- HrECHR: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1 *w/w*) of *E. coccinea* + *H. rosasinensis* (20 mg/kg in PBS; i.p.).
- HrAGEH: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1:1 *w/w*) of *A. gangetica* + *E. coccinea* + *H. rosasinensis* (20 mg/kg in PBS; i.p.).
- HrAMAE: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1:1 *w/w*) of *A. montanus* + *A. gangetica* + *E. coccinea* (20 mg/kg in PBS; i.p.).
- HrAMAH: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1:1 *w/w*) of *A. montanus* + *A. gangetica* + *H. rosasinensis* (20 mg/kg in PBS; i.p.).
- HrAMEH: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1:1 *w/w*) of *A. montanus* + *E. coccinea* + *H. rosasinensis* (20 mg/kg in PBS; i.p.).
- HrAAEH: HyGR received SCF + water *ad libitum* + combined dose (ratio: 1:1:1:1 *w/w*) of *A. montanus* + *A. gangetica* + *E. coccinea* + *H. rosasinensis* (20 mg/kg in PBS; i.p.).

At the end of treatment, the animals were fasted for 12 h<sup>28</sup> before their blood samples and organ homogenates were measured for the various biochemical parameters.

### Fasting plasma glucose concentration

Blood samples (2.0 mL) were drawn from the orbital sinus<sup>42</sup> on the 30th day and measured for FPGC. Determination of FPGC was by the glucose oxidase method according to Randox® kit manufacturer's procedure (Randox® Laboratories Ltd. Ardmore, United Kingdom).

### Preparation of renal and hepatic organs homogenates

The rats were anaesthetized in CCl<sub>4</sub> chamber and dissected. The kidneys and liver were quickly excised and placed on a blotting paper to remove blood and rinsed in 1.15% of potassium chloride solution to remove residual hemoglobin.<sup>43</sup> The two organs were stored in 10% formaldehyde and preserved at temperature below -8°C before analyses.<sup>43</sup> with minor modifications by Chikezie and Uwakwe.<sup>44</sup> Each organ was homogenized using a Teflon homogenizer in aqueous K<sub>2</sub>PO<sub>4</sub>/KHPO<sub>4</sub> buffer (0.1 M; pH=7.4); in 4:1 volume of buffer to organ weight. Next, the crude homogenate was centrifuged at 10,000 ×g for 20

min at 4°C to obtain the post mitochondrial supernatant fraction (PMSF). The PMSF was finally stored at -8 °C before used for analyses. The PMSF was measured for GPOx, SOD and CAT activities. Protein concentrations of PMSF of hepatic and renal homogenates were assayed<sup>45</sup> using bovine serum albumin (Sigma-Aldrich, St. Louis, USA) as a standard. Also, the PMSF was measured for glutathione (GSH) concentration.

#### Glutathione peroxidase activity

GPOx activity was measured by the method of Paglia and Valentine<sup>46</sup> as reported by Chikezie and Uwakwe.<sup>44</sup> Briefly, the reaction mixture contained aliquot of PMSF in 50 mM K<sub>2</sub>PO<sub>4</sub>/KHPO<sub>4</sub> buffer (pH=7.0), 1.0 mM EDTA (Mayer and Baker, England), 1.0 mM NaN<sub>3</sub>, 0.2 mM NADPH, 1.0 U glutathione reductase, and 1.0 mM GSH. The reaction mixture was allowed to equilibrate at 25°C for 5 min. The reaction was started by introducing 0.1 mL of 2.5 mM H<sub>2</sub>O<sub>2</sub>. Increase in absorbance at λ<sub>max</sub>=340 nm was recorded for 5 min. The change in absorbance was defined as nanomoles of NADPH oxidized to NADP; Σ=6.2 × 10<sup>3</sup> M<sup>-1</sup> cm<sup>-1</sup> at λ<sub>max</sub>=340 nm. The levels of GPx were expressed in terms of nmole NADPH consumed/min/mg protein (U/mg protein).

#### Superoxide dismutase activity

SOD activity was measured.<sup>47</sup> Briefly, the reaction mixture contained solution A (50 mM Na<sub>2</sub>CO<sub>3</sub>, 0.1 mM EDTA, pH=10.0), solution B (96 μM nitroblue tetrazolium [NBT] in solution A), and solution C (0.6% Triton X-100 in solution A) were incubated at 37°C for 10 min. Reaction was started by introducing 100 μL of solution D (20 mM hydroxylamine hydrochloride, pH=6.0). The rate of NBT dye reduction by O<sub>2</sub><sup>-</sup> anion generated due to photo-activation of hydroxylamine hydrochloride was measured at λ<sub>max</sub>=560 nm in the absence of PMSF. Next, 10 μL aliquot of PMSF were added to the reaction mixture and 50% inhibition in the rate of NBT reduction by SOD present in the enzyme source was measured. A unit (U) of SOD activity was defined by the 50% inhibition of NBT. SOD activity was expressed in U/mg protein.

#### Catalase activity

Measurement of PMSF CAT activity was according to the method of Luck,<sup>48</sup> as reported by Chikezie and Uwakwe.<sup>44</sup> The final reaction volume of 3.0 mL contained 0.05 M Tris-buffer, 5 mM EDTA (pH=7.0), and 10 mM H<sub>2</sub>O<sub>2</sub> (in 0.1 M K<sub>2</sub>PO<sub>4</sub>/KHPO<sub>4</sub> buffer; pH=7.0). A hundred micro milliliter (100 μL) aliquot of the PMSF was added to the above mixture. The rate of change of absorbance per min at λ<sub>max</sub>=240 nm was recorded for 5 min. CAT activity was

calculated using Σ=43.6 M<sup>-1</sup>cm<sup>-1</sup> and expressed in terms of mole H<sub>2</sub>O<sub>2</sub> consumed/min/mg protein (U/mg protein).

#### Reduced glutathione (GSH)/oxidized glutathione (GSSG) ratio

Level of GSH in organ homogenate was determined<sup>49</sup> with minor modification by Chikezie and Uwakwe.<sup>44</sup> A 100 μL aliquot of the PMSF was mixed with 25% of CHCl<sub>3</sub> and centrifuged at 2000 × g for 15 min to precipitate proteins. The supernatant was aspirated and diluted to 1.0 mL with 0.2 M Na<sub>2</sub>PO<sub>4</sub>/NaHPO<sub>4</sub> buffer (pH=8.0). Next, 2.0 mL of 0.6 mM 5, 5'-dithiobis-(2-nitrobenzoic acid) (DTNB) was added. The absorbance of the developed yellow-colour complex maintained at 24±5°C was measured at λ<sub>max</sub>=405 nm after 10 min. A standard curve was obtained with GSH standards. The level of GSH was expressed as μg GSH/mg protein. The GSSG concentrations (C<sub>GSSG</sub>) of organ homogenates were measured using Bioxytech-412 kits according to manufacturer's procedure (Oxis International Inc., Foster City, CA, USA).

$$C_{GSSG} = \frac{Abs_{blank} - Abs_{sample} \times C_{std}}{Abs_{blank} - Abs_{std}}$$

Accordingly, the ratio of GSH/GSSG of renal and hepatic tissues was evaluated.

#### Statistical analysis

The data collected were analyzed by the analysis of variance procedure while treatment means were separated by the least significance difference (LSD) incorporated in the statistical analysis system (SAS) package of 9.1 version, (2006). Correlation coefficients and trend lines were determined using Excel Software (Microsoft, 2010 version).

## RESULTS

At the end of the experimental time of 30 days, FPGC of DIAB group was 4.3 folds > NORM group (Table 1). Specifically, HrACMO, HrHIRO, HrAGHR and HrAMEH groups exhibited elevated FBGC compared with the NORM group (*p*<0.05) and were considered to be hyperglycemic after the 30-day treatment. Conversely, FPGC of HrASGA, HrAMEC, HrAMHR, HrECHR, HrAGEH and HrAAEH groups were significantly (*p*<0.05) higher than the NORM group but not considered to be hyperglycemic after the 30-day treatment.

It is worthwhile to note that FPGC of HrAGAM and HrAGEC groups were significantly (*p*<0.05) lower than the NORM group. Specifically, HrAGAM group=66.30

**Table 1: Fasting plasma glucose concentrations of normal, diabetic and treated rats**

Group	FPGC (mg/dL)
NORM	86.30 ± 0.15 <sup>n</sup>
DIAB	368.20 ± 0.72 <sup>a</sup>
HrACMO	133.00 ± 0.63 <sup>b</sup>
HrASGA	105.70 ± 0.71 <sup>e,f,g,h</sup>
HrEMCO	118.00 ± 0.92 <sup>c,d</sup>
HrHIRO	112.50 ± 0.70 <sup>d,e</sup>
HrAGAM	66.30 ± 0.71 <sup>q</sup>
HrAGEC	81.00 ± 0.39 <sup>n,o,p</sup>
HrAGHR	112.50 ± 0.42 <sup>d,e,f</sup>
HrAMEC	97.20 ± 0.51 <sup>l,m</sup>
HrAMHR	105.00 ± 0.62 <sup>e,f,g,h,i</sup>
HrECHR	103.50 ± 0.79 <sup>f,g,h,i,j</sup>
HrAGEH	103.00 ± 0.81 <sup>f,g,h,i,j,k</sup>
HrAMAE	84.70 ± 0.51 <sup>n,o</sup>
HrAMAH	124.30 ± 0.92 <sup>c</sup>
HrAMEH	112.00 ± 0.71 <sup>d,e,f,g</sup>
HrAAEH	97.30 ± 0.85 <sup>l</sup>

The mean (X) ± S.D of six (n = 6) determinations. Means in the column with the same letter are not significantly different at  $p > 0.05$  according to LSD. FPGC > 110 mg/dL = hyperglycemia.

± 0.71 mg/dL, HrAGEC group=81.00±0.39 mg/dL and NORM group=86.30±0.15 mg/dL (Table 1). Generally, FPGC of the various treated HyGR was within the range of 66.30±0.71 mg/dL and 133.00±0.63 mg/dL as against DIAB group=368.20±0.72 mg/dL. At the end of the experimental time, the following animal groups: HrASGA, HrAGAM, HrAGEC, HrAMEC, HrAMHR, HrECHR, HrAGEH, HrAMAE and HrAAEH registered FPGC < 110 mg/dL.

Figure 1 shows that flavonoids were comparatively the most abundant phytochemical present in the four leaves extracts, in the order: AMCO > ASGA > HIRO > EMCO. The concentration of alkaloids in HIRO was relatively low compared with other three leaf extracts. Saponin contents in the four leaf extracts were in the range of 43.50±1.01 mg/g and 71.01±0.62 mg/g. In addition, saponin contents amongst ASGA, EMCO and HIRO leaf extracts were not significantly ( $p > 0.05$ ); specifically, ASGA=44.50±0.51 mg/g, EMCO=43.50±0.54 mg/g and HIRO=45.25±0.35 mg/g (Figure 1). Likewise, tannin contents of the four leaf extracts was within a narrow range of 25.50±0.18 mg/g and 33.75 ± 0.32 mg/g;  $p > 0.05$ . In addition, tannin was the lowest phytochemical present in the four leaf extracts.


Liver SOD (LSOD) activity of NORM group was significantly ( $p < 0.05$ ) higher the corresponding kidney SOD (KSOD) activity (Figure 2). Likewise, in the DIAB group,

LSOD activity was higher than KSOD activity;  $p > 0.05$ . Also, Figure 2 shows that the levels of LSOD and KSOD activities of the DIAB group were in 3.82 and 12.01 folds lower than the NORM group, respectively. At the end of the experimental time, each of the treated diabetic rats LSOD activities were lower than that of NORM group; except HrAMHR<sub>LSOD</sub> activity=198.24±1.02 U/mg protein, whereas NORM<sub>LSOD</sub> activity=190.61±1.22 U/mg protein;  $p > 0.05$ . By the same comparison, HrAGAM<sub>KSOD</sub> activity and HrAGHR<sub>KSOD</sub> activity were significantly ( $p < 0.05$ ) higher than NORM<sub>KSOD</sub> activity. Peak values of SOD activities of the treated HyGR were of the following magnitudes: HrACMO<sub>LSOD</sub> activity=185.25±1.01 U/mg proteins and HrAGAM<sub>KSOD</sub> activity=171.45±0.99 U/mg protein.


However, KSOD activities of other treated HyGR (HrAGEC, HrAMAH, HrECHR and HrAGEH) were higher than the NORM group, but with no significant difference ( $p > 0.05$ ). Generally, the treated HyGR exhibited increased levels of LSOD and KSOD activities compared with the DIAB group. For instance, HrAMHR group showed 3.97 folds improvement in LSOD activity compared with DIAB group. LSOD and KSOD activities of the treated HyGR showed a very weak positive correlation ( $r = 0.197011749$ ).

Liver GPOx (LGPOx) activity of the NORM group was not significantly ( $p > 0.05$ ) different from the corresponding kidney GPOx (KGPOx) activity; LGPOx activity=1.70±0.03 U/mg protein, whereas KGPOx activity=1.10 ± 0.04 U/mg protein (Figure 3). LGPOx and KGPOx activities of DIAB group were 2.36 folds and 3.93 folds lower than the NORM group. However, LGPOx and KGPOx activities increased following the administration of the various herbal formulations to corresponding animal groups as against the decreased levels of LGPOx and KGPOx activities of DIAB group. LGPOx activity showed a very weak negative correlation ( $r = -0.088487792$ ) with KGPOx activity in the treated animal groups. Figure 3 shows that amongst the various treated hyperglycemic rat groups, peak LGPOx activity=1.71 ± 0.05 U/mg proteins, occurred following the administration of AAEH herbal formulation. Similarly, peak value of KGPOx activity=1.99±0.04 U/mg protein, ensued in HyGR consequent upon the administration of AMHR herbal formulation.


Liver CAT (LCAT) activity was 2.19 folds > kidney CAT (KCAT) activity in NORM group (Figure 4). DIAB<sub>LCAT</sub> activity=23.05±0.82 U/mg protein and DIAB<sub>KCAT</sub> activity=20.78±0.77 U/mg protein, representing 73.93% and 48.60% reduction in CAT activity compared with NORM



**Figure 1:** Some phytochemical contents of *A. montanus*, *A. gangetica*, *E. coccinea* and *H. rosasinensis* leaf extracts.




**Figure 2:** Liver and kidney superoxide dismutase activities of experimental rats.




**Figure 3:** Liver and kidney glutathione peroxidase activities of experimental rats.

group;  $p < 0.05$ . Comparative analyses showed that  $DIAB_{LCAT}$  activity was not significantly ( $p > 0.05$ ) different from  $DIAB_{KCAT}$  activity. Furthermore, at the end of the experimental time, LCAT activity and KCAT activity of all the treated hyperglycemic rat groups were significantly ( $p < 0.05$ ) lower than the NORM group; except  $HrASGA_{KCAT}$  and  $HrEMCO_{KCAT}$  activities;  $p > 0.05$ . LCAT activity of the treated hyperglycemic animal groups showed a very weak positive correlation ( $r = 0.000829808$ ) with that of KCAT activity. HrAAEH group exhibited peak LCAT activity =  $83.77 \pm 0.82$  U/mg protein, whereas HrEMCO group gave peak KCAT activity =  $41.22 \pm 0.72$  U/mg protein.



**Figure 4:** Liver and kidney catalase activities of experimental rats.

The ratio of liver GSH/GSSG (LGSH: GSSG) levels and kidney GSH/GSSG (KGSH: GSSG) levels were such that in NORM group,  $KGSH: GSSG > LGSH: GSSG$ ;  $p < 0.05$  (Figure 5). Also, Figure 4 shows that most values of liver and kidney GSH/GSSG ratios were significantly ( $p < 0.05$ ) increased when compared with NORM group, following the administration of the herbal formulations to the various hyperglycemic rat groups; except HrAMAE group which gave  $LGSH: GSSG = 9.74 \pm 1.04$  and  $KGSH: GSSG = 19.00 \pm 1.06$ . LGSH: GSSG and KGSH: GSSG of treated hyperglycemic rat groups showed a very weak negative correlation ( $r = -0.192889137$ ). Fig-



**Figure 5:** Liver and kidney ratio of GSH to GSSG concentrations of experimental rats.

Figure 5 shows that LGSH: GSSG peak value =  $15.74 \pm 1.05$ , occurred in treated hyperglycemic rat groups administered with AMHR herbal formulation. Correspondingly, peak value of KGSH: GSSG =  $26.11 \pm 1.04$ , was presented in HyGR administered with AMEC herbal formulation.

An overview of Figures 2-5 showed that liver and kidney levels of antioxidant enzyme activities (SOD, GPOx and CAT) and GSH/GSSG ratios increased following the administration of the various herbal formulations, which was order of SHF > DHF > THF > QHF as indicated by their corresponding trend lines.

## DISCUSSION

From the results of the present study, HyGR treated with the various herbal formulations for experimental period of 30 days showed evidence of varied levels of efficacy of herbal formulations to exert glycemic control (Table 1). Specifically, HyGR administered with the DHFs (HrAGAM, HrAGEC and HrAMEC), THF (HrAMAE) and QHF (HrAAEH) were normoglycemic (FBGC =  $66.30 \pm 0.71 - 97.30 \pm 0.85$  mg/dL) following the 30-day treatments. In conformity with the present findings, several authors had previously reported the use of herbal remedies for the control and treatment of hyperglycemia in animal models<sup>28,32,50,51</sup> and individuals suffering from diabetes mellitus.<sup>52-56</sup> Furthermore, the modes of actions of some selected anti-hyperglycemic plants have been extensively discussed elsewhere.<sup>32,57,58</sup> In all those reports, the quantity and types of phytochemicals present in the extracts were considered to be, to a large extent, responsible for the observed anti-hyperglycemic properties of the herbal extracts.<sup>32,50,51,53,56,59,60</sup>

Previous studies showed that alkaloids and flavonoids are effective anti-diabetic agents.<sup>57,61</sup> In another study, the potency of core structures of flavonoid precursors to act as anti-hyperglycemic agents was demonstrated *in vivo* in HyGR.<sup>62</sup> A comprehensive review on natural approach to the treatment of diabetes,<sup>7</sup> further confirmed that flavonoids are among the listed anti-diabetic compounds, which exert their hypoglycemic properties via extra pancreatic mechanism of  $\alpha$ -glucosidase modulation.<sup>56</sup> Expectedly, the relatively high flavonoids contents of the four medicinal plants (ACMO, ASGA, EMCO and HIRO) contributed to the capabilities of the herbal formulations to exert glycemic control in the experimental rats. Likewise, the relatively high abundance of alkaloids, especially in ACMO, ASGA and EMCO extracts, may have acted as a hypoglycemic agent as previously reported.<sup>63</sup> Therefore, combinatorial formulations of the four medicinal plant extracts promoted synergy amongst the various bioactive principles, and thereby potentiated the anti-hyperglycemic capacities of the individual plant extracts in conformity with previous reports.<sup>33,64</sup> Similarly, Brahmabhatt *et al.*,<sup>65</sup> demonstrated that combining ginger extracts' with other constituents, particularly 6-gingerol, caused significant augmentation of anti-proliferative activity of the extracts. However, studies have also shown that antagonistic interactions among phytochemicals in medicinal plant extracts could, paradoxically, attenuate the therapeutic potentials of the individual bioactive principles.<sup>66</sup>

Saponins and tannins are bioactive principles of medicinal and toxicity importance and their capabilities to exert glycemic control have been experimentally established.<sup>67-69</sup> Kunyanga *et al.*,<sup>67</sup> reported that condensed tannin extracts of raw and processed indigenous food ingredients from Kenya exhibited promising anti-diabetic effects; possessing potential  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibition activ-



ities within the range of 23% to 44% and 58% to 88%, respectively. In another study Zheng *et al.*,<sup>68</sup> implicated the saponins as the anti-diabetic principles of seed extract of *Entada phaseoloides* L. They further posited that the therapeutic effect of saponins was facilitated by repression of chronic inflammation response pathways of pancreatic islets that was mediated by its inhibitory actions on interleukin-6 (IL-6), tumor necrosis factor-alpha (TNF- $\alpha$ ) and C-reactive protein (CRP) by the seed extracts in a dose dependent manner. Histopathological examination of the pancreatic tissues of streptozotocin-induced diabetic rats (S-IDR) showed evidence of considerable quantitative increase in  $\beta$ -cells by 75% when treated with the saponins.<sup>69</sup> The present study showed that saponins and tannins are present in approximately equal proportions in extracts of ACOMO, ASGA, EMCO and HIRO that may have contributed, in part, to the anti-hyperglycemic potentials of the experimental plant extracts.

Generally, the level of enzyme activity in biologic systems could serve as a reliable tool in ascertaining health and pathologic conditions.<sup>20</sup> Also, in the event of exposure of biological systems to noxious chemical agents and other environmental insults, previous researchers have applied levels of activity of diverse antioxidant enzymes and low molecular weight antioxidants (LMWA) as basis for ascertaining tissue oxidative stress status.<sup>18,20,26,70-73</sup> Findings from the present study showed that HyGR exhibited alterations in tissue antioxidant enzymes (SOD, GPOx and CAT) activities and GSH/GSSG ratio, of which the cellular levels these oxidative stress indicators were diagnostic of oxidative tissue injury.

Decreased SOD activity of untreated HyGR as reported here conformed with previous studies.<sup>11,74,75</sup> In particular, Di Naso *et al.*,<sup>11</sup> noted that decreased SOD activity in DM is connected with non-enzymatic glycation of intracellular SOD at the lysine residues located in the heparin-binding domain, which indirectly altered extracellular SOD activity and functionality. In another study Loven *et al.*,<sup>76</sup> posited that decreased Cu-Zn SOD activity observed in diabetes was sequel to the inactivation of the enzyme as a result of absence or failure to metabolize hydrogen peroxide via GPOx pathway. The decreased levels of renal and hepatic SOD activity of HyGR showed evidence of restoration of the enzyme activity that were comparable to NORM rats following the administration of the various experimental herbal formulations (Figure 2). The present findings are comparable with that reported by Elmalí *et al.*,<sup>77</sup> in which they noted that glibenclamide directly increased renal and hepatic CAT and SOD activities of S-IDR. Reports have shown that ethanolic extract of

*Sphaeranthus indicus* exhibited protective effect against lipid peroxidation and normalized repressed SOD, CAT, glutathione S-transferase activities in gentamicin induced nephrotoxic rats.<sup>70</sup> Similarly, previous reports showed that S-IDR treated with exogenous SOD and GSH exhibited improved antioxidant enzymes activities,<sup>11,74,76</sup> which implied amelioration of tissue oxidative stress.

Depressed renal and hepatic GPOx and CAT activities are associated with the diabetic state as reported here (Figures 3 and 4). Specifically, previous studies<sup>78</sup> had noted that CAT deficiency accelerated diabetic nephropathy, which was mediated by peroxisomal dysfunction. However, cod liver oil intake engenders significant increase in both GPOx and CAT activities in aorta, heart, and liver of S-IDR,<sup>79</sup> thereby alleviating oxidative stress induced tissue injuries. Similarly, studies by Shanmugam *et al.*,<sup>75</sup> revealed that dietary ginger improved renal and hepatic GPOx and CAT activities, alongside other antioxidant enzymes and oxidative stress indicators of S-IDR. They further noted that treatment of diabetic rats with ginger for 30 days caused a therapeutic protective effect by lowering oxidative stress induced hepatic and renal damage. Also, using experimental female Wistar rats, several natural products obtained from wide varieties of medicinal plants reversed reduced tissue SOD, CAT, GPOx and GST activities following exposure to cadmium.<sup>72</sup> They further suggested that their therapeutic actions were probably connected with the synergistic effects of the bioactive principles of the various herbal extracts. The present study has equally shown substantial improvements in depressed renal and hepatic GPOx and CAT activities of HyGR following the administration of combinatorial herbal formulations of the four medicinal plant extracts.

The physiochemical roles of GSH is indispensable in a multitude of cellular processes. Therefore, disturbances in GSH homeostasis are associated with the etiology and/or progression of a number of human diseases.<sup>26</sup> Furthermore, the redox state of the GSH/glutathione disulfide couple (GSH/GSSG) is an important biologic indicator and regulator of signals of and metabolic pathways in normal cell systems<sup>27,80</sup> as well as oxidative stress and glucose metabolism in type 2 DM.<sup>81</sup> Accordingly, the results of the present investigations have shown that GSH/GSSG ratio served as a reliable biomarker in ascertaining the efficacy of herbal remedies in alleviation of hyperglycemia and amelioration of tissue oxidative stress. The present findings corroborated the reports of El-Missiry and El Gindy,<sup>82</sup> in which it was noted that daily oral administration of *Eruca sativa* seeds oil for 2 weeks stimulated and increased hepatic GSH production of

alloxan-treated rats. Additionally, Zitka *et al.*,<sup>83</sup> used the redox status expressed as GSH:GSSG as a biomarker for ascertaining the level of oxidative stress in paediatric tumor patients, which was similar to earlier reports<sup>84</sup> on impairment of glutathione metabolism in erythrocytes of patients suffering from diabetes mellitus. From the present investigations, the antioxidant enzymes activities and GSH:GSSG between renal and hepatic tissues of HyGR treated with the various herbal formulations gave poor correlations. This was an obvious indication that the levels of oxidative stress and subsequent capacities of the herbal formulations to ameliorate oxidative stress in renal and hepatic tissues of HyGR were not of equal pathophysiological proportions and dimensions.

## CONCLUSION

Generally, the improvement in renal and hepatic antioxidant enzymes activities and GSH:GSSG of the treated

HyGR suggest that the experimental herbal formulations caused a sparing effect on renal and hepatic antioxidant system against reactive oxygen species. Additionally, the combinatorial herbal formulations exerted greater glycaemic control than the single herbal formulations.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ACKNOWLEDGEMENT

The authors are thankful to Dr. M. Ibe, School of Agriculture and Agricultural Technology (SAAT), Federal University of Technology, Owerri.

## REFERENCES

- Adeyemi AO, Okpo SO, Okpaka O. The analgesic effect of the methanolic extract of *Acanthus montanus*. *J Ethnopharmacol.* 2004; 90(1): 45-8.
- Okoli CO, Akah PA, Onuoha NJ, Okoye TC, Nwoye AC, Nworu CS. *Acanthus montanus*: An experimental evaluation of the antimicrobial, anti-inflammatory and immunological properties of a traditional remedy for furuncles. *BMC Compl Altern Med.* 2008; 8(1): 27. doi: 10.1186/1472-6882-8-27.
- Ruffo CK, Birnie A, Tengas B. Edible wild plants of Tanzania. *Relma Technical Handbook*; 2002. P. 766.
- Kokwaro J: *Medicinal plants of East Africa.* 2<sup>nd</sup> Ed. Kenya Lit Bureau. Nairobi, Kenya; 1993. P. 401.
- Siddiqui AA, Wani SM, Rajesh R, Alagarsamy V. Phytochemical and pharmacological investigation of flowers of *Hibiscus rosasinensis* linn. *Indian J Pharm Sci.* 2006; 68(1): 127-30.
- Soni D, Gupta A, Solanki R, Jana GK. Pharmacognostical, phytochemical and physicochemical findings over the root extract of *Hibiscus rosasinensis* [Malvaceae]. *J Nat Prod Plant Resour.* 2011; 1(4): 73-9.
- Kumar S, Narwal S, Kumar V, Prakash O.  $\alpha$ -Glucosidase inhibitors from plants: a natural approach to treat diabetes. *Pharmacog Rev.* 2011; 5(9): 19-29.
- Argano M, Brignardello E, Gatto VTO, Danni O, Boccuzzi G. Dehydroepiandrosterone administration prevents the oxidative damage induced by acute hyperglycemia in rats. *J Endocrinol.* 1997; 155(2): 233-40.
- Hammes H-P, Du X, Edelstein D, Taguchi T, Matsumura T, Ju Q, *et al.* Benfotiamine blocks three major pathways of hyperglycemic damage and prevents experimental diabetic retinopathy. *Nat Med.* 2003; 9(3): 294-9.
- Evans JL, Goldfine ID, Maddux BA, Grodsky GM. Oxidative stress and stress activated signaling pathways: a unifying hypothesis of type 2 diabetes. *Endocr Rev.* 2002; 23(5): 599-622.
- Di Naso FC, Dias AS, Porawski M, Marroni NAP. Exogenous superoxide dismutase: action on liver oxidative stress in animals with streptozotocin-induced diabetes. *Exp Diabetes Res.* 2011; 2011: 754132. 7pages. doi: 10.1155/2011/754132
- Vaag A, Dmsbo P, Hother-Nielsen O, Beck-Nielsen H. Hypoglycemia compensates for the defects in insulin mediated glucose metabolism and in the activation of glycogen synthase in the skeletal muscle of patients with type 2 (non-insulin dependent) diabetes mellitus. *Diabetologia* 1992; 35: 80-8.
- Babaei-Jadidi R, Karachalias N, Ahmed N, Battah S, Thornalley PJ. Prevention of incipient diabetic nephropathy by high-dose thiamine and benfotiamine. *Diabetes* 2003; 52(8): 2110-20.
- Williamson JR, Chang K, Frangos M, Hasan KS, Ido Y, Kawamura T. Hyperglycemia pseudo hypoxia and diabetic complications. *Diabetes* 1993; 42(6): 801-13.
- Wolff SP, Jiang ZY, Hunt JV. Protein glycation and oxidative stress in diabetes mellitus and ageing. *Free Radic Biol Med.* 1991; 10(5): 339-52.
- Ceriello A, Quattraro A, Giugliano D. New insight on non-enzymatic glycosylation may lead to therapeutic approaches for the prevention of the diabetic complications. *Diabetes Med.* 1992; 9(3): 297-9.
- Ohtsuka Y, Agishi Y. Abnormal amino acid metabolism in diabetes mellitus. *Nihon Rinsho.* 1992; 50(7): 1631-6.
- Mate's JM. Effects of antioxidant enzymes in the molecular control of reactive oxygen species toxicology. *Toxicol.* 2000; 153(1): 83-104.
- Bonnefont-Rousselot D, Bastard JP, Jaudon MC, Delattre J. Consequences of diabetic status on the oxidant/antioxidant balance. *Diabetes Med.* 2000; 26(3): 163-76.
- Kohen R, Nyska A. Oxidation of biological systems: oxidative stress phenomena, antioxidants, redox reactions, and methods for their quantification. *Toxicol Pathol.* 2002; 30(6): 620-50.
- Imai H, Narashima K, Arai M, Sakamoto H, Chiba N, Nakagawa Y. Suppression of leukotriene formation in RBL-2H3 cells that overexpressed phospholipid hydroperoxide glutathione peroxidase. *J Biol Chem.* 1998; 273(4): 1990-7.
- Guan Y, Hickey MJ, Borgstahl GE, Hallewell RA, Lepock JR, O'Connor D, *et al.* Crystal structure of Y34F mutant human mitochondrial manganese superoxide dismutase and the functional role of tyrosine 34. *Biochem.* 1998; 37(14): 4722-30.
- Banci L, Benedetto M, Bertini I, Del Conte R, Piccoli M, Viezzoli MS. Solution structure of reduced monomeric Q133M2 copper, zinc superoxide dismutase (SOD). Why is SOD a dimeric enzyme? *Biochem.* 1998; 37(34): 11780-91.
- Young H, Kim EJ, Roe JH, Hah YC, Kang SO. A novel nickel-containing superoxide dismutase from *Streptomyces spp.* *Biochem J.* 1996; 318: 889-96.
- Schafer FQ, Buettner GR. Redox environment of the cell as viewed through the redox state of the glutathione disulfide/glutathione couple. *Free Radic Biol Med.* 2001; 30(11): 1191-212.
- Ballatori N, Krance SM, Notenboom S, Shi S, Tieu K, Hammond CL. Glutathione dysregulation and the etiology and progression of human diseases. *Biol Chem.* 2009; 390(3): 191-214.
- Kemp M, Go YM, Jones DP. Nonequilibrium thermodynamics of thiol/disulfide redox systems: a perspective on redox systems biology. *Free Radic Biol Med.* 2008; 44(6): 921-37.

28. El-Demerdash FM, Yousef MI, Abou El-Naga NI. Biochemical study on the hypoglycemic effects of onion and garlic in alloxan-induced diabetic rats. *Food Chem Toxicol.* 2005; 43(1): 57–63.
29. Anwar F, Latif S, Ashraf M, Gilani AH. *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytother Res.* 2007; 21(1): 17–25.
30. Gao D. Antioxidant Therapies for Hypolipidemia and Hyperglycemia; 2012. <http://dx.doi.org/10.5772/47432>. [Last retrieved on 2014 June 09].
31. Mungle AN, Bodhankar NM, Chandak KK. Antidiabetic potential of *Dolichandrone falcata* leaves in alloxan induced diabetic rats. *Int J Res Pharmaceut Biomed Sci.* 2012; 3(1): 319-24.
32. Patel DK, Prasad SK, Kumar R, Hemalatha S. An overview on antidiabetic medicinal plants having insulin mimetic property. *Asian Pac J Trop Biomed.* 2012; 2(4): 320–30.
33. Wang F, Zhao S, Li F, Zhang B, Qu Y, Sun T, *et al.* Investigation of antioxidant interactions between *Radix Astragali* and *Cimicifuga foetida* and identification of synergistic antioxidant compounds. *Plos One.* 2014; 9(1): e87221. doi:10.1371/journal.pone.00872.
34. Matteucci E, Giampietro O. Oxidative stress in families of type 1 diabetic patients. *Diabetes Care* 2000; 23(8): 1182-6.
35. Brownlee M. Biochemistry and molecular cell biology of diabetic complications. *Nature* 2001; 414(6865): 813–20.
36. Boham AB, Kocipai AC. Flavonoid and condensed tannins from leaves of Hawaiian *Vaccinium vaticulum* and *V. calycinium*. *Pacific Sci.* 1994; 48: 458-63.
37. Harborne JB. *Phytochemical methods*, London. Chapman and Hall, Ltd; 1973. p. 49-188.
38. Obadoni BO, Ochuko PO. Phytochemical studies and comparative efficacy of crude extracts of some homeostatic plants in Edo and Delta States of Nigeria. *Global J Pure Appl Sci.* 2001; 8: 203-8.
39. Van-Burden TP, Robinson WC. Formation of complexes between protein and tannin acid. *J Agric Food Chem.* 1981; 1(4): 77-82.
40. Belonwu DC, Ibegbulem CO, Nwokocho MN, Chikezie PC. Some phytochemicals and hydrophilic vitamins of *Anacardium occidentale*. *Res J Phytochem.* 2014; 8(3): 78-91.
41. Ibegbulem CO, Chikezie PC. Hypoglycemic properties of ethanolic extracts of *Gongronema latifolium*, *Aloe perryi*, *Viscum album* and *Allium sativum* administered to alloxan-induced diabetic albino rats (*Rattus norvegicus*). *Pharmacog Commun.* 2013; 3(2): 12-6.
42. Hoff N. Methods of blood collection in the mouse. *Lab Ani.* 2000; 29(10): 47-53.
43. Adekunle AS, Adedeji AL, Oyewo EO, Adedosu OT, Omotoso AT. Hyperlipidemia induced by atherogenic diet enhanced oxidative stress in the kidney and inflammatory responses: an *in-vivo* study. *Asian J Nat Appl Sci.* 2013; 2(1): 82-93.
44. Chikezie PC, Uwakwe AA. Activities of three erythrocyte enzymes of hyperglycemic rats (*Rattus norvegicus*) treated with *Allium sativa* extract. *J Diabetes Metab Disorders.* 2014; 13(1): 50, 9 pages. doi: 10.1186/2251-6581-13-50.
45. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the folin phenol reagent. *J Biol Chem.* 1951; 13(1): 265–75.
46. Paglia DE, Valentine WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med.* 1967; 70(1): 158–69.
47. Kono Y. Generation of superoxide radical during autoxidation of hydroxylamine and an assay for superoxide dismutase. *Arch Biochem Biophys.* 1978; 186(1): 189–95.
48. Luck H. Catalase. In: *Methods of enzymatic analysis* (Bergmeyer H.W. Ed.). Academic Press, New York. 1963; Section 3: 885–94.
49. Moron MS, Kepierre JW, Mannervick B. Levels of glutathione reductase and glutathione S-transferase activities in rat lung and liver. *Biochim Biophys Acta.* 1979; 582(1): 67–8.
50. Pierre W, Gildas AJH, Ulrich MC, Modeste W-N, Benoît NT, Albert K. Hypoglycemic and hypolipidemic effects of *Bersama engleriana* leaves in nicotinamide/streptozotocin-induced type 2 diabetic rats. *BMC Compl Altern Med.* 2012; 12(1): 264. 6pages.
51. Onunogbo CC, Ohaeri OC, Eleazu CO. Effect of mistletoe (*Viscum album*) extract on the blood glucose, liver enzymes and electrolyte balance in alloxan induced diabetic rats. *Am J Bio chem MolBiol.* 2013; 3: 143–50.
52. Gray AM, Flatt PM. Insulin-releasing and insulin-like activity of the traditional anti-diabetic plant *Coriandrum sativum* (coriander). *Br J Nutri.* 1999; 81(03): 203-9.
53. Harmanto N. Conquering disease in unison with Mahkota Dewa, In: Harmanto (Ed.), *PT Mahkota Dewa Indonesia*, North Jakarta; 2003. p.14.
54. Jayawardena MHS, de Alwis NMW, Hettigoda V, Fernando DJS. A double blind randomized placebo controlled cross over study of herbal preparation containing *Salacia reticulata* in the treatment of type II diabetes. *J Ethnopharmacol* 2005; 96: 215–8.
55. Ahmed MF, Kazim SM, Ghorri SS, Mehjabeen SS, Ahmed SR, Ali SM, *et al.* Antidiabetic activity of *Vincarosea* extracts in alloxan-induced diabetic rats. *Int J Endocrinol.* 2010; Article ID 841090: 6 pages. doi:10.1155/2010/841090.
56. Ali RB, Atangwho IJ, Kuar N, Mohamed EAH, Mohamed AJ, Asmawi MZ, *et al.* Hypoglycemic and anti-hyperglycemic study of *Phaleria macrocarpa* fruits pericarp. *J Med Plants Res.* 2012; 6(10): 1982–90.
57. Jung M, Park M, Lee HC, Kang YH, Kang ES, Kim SK. Anti-diabetic agents from medicinal plants. *Curr Med Chem.* 2006; 13(10): 1203–18.
58. Prabhakar PK, Doble M. A target based therapeutic approach towards diabetes mellitus using medicinal plants. *Curr Diabetic Rev.* 2008; 4(4): 291-308.
59. Daisy P, Priya CE, Vargese L. A study on the regenerative potential of the root and leaf extracts of *Elephantopus scaber* L.: An antidiabetic approach. *Afr J Pharm Pharmacol.* 2011; 5(16): 1832-7.
60. Ojiako AO, Chikezie PC. Comparative proximate composition and hypoglycemic properties of three medicinal plants (*Veronia amygdalina*, *Azadirachta indica* and *Moringaoleifera*). *Pharmacog Commun.* 2014; 4(3): 40-8.
61. Chen F, Xiong H, Wang J, Ding X, Shu G, Mei Z. Antidiabetic effect of total flavonoids from *Sanguis draxonis* in type 2 diabetic rats. *J Ethnopharmacol.* 2013; 149(3): 729-36.
62. Najafian M, Ebrahim-Habibi A, Yaghmaei P, Parivar K, Larijani B. Core structure of flavonoids precursor as an anti-hyperglycemic and anti-hyperlipidemic agent: An *in vivo* study in rats. *Acta Biochim Pol.* 2010; 57(4): 553-60.
63. Saidu AN, Mann A, Balogun S. The hypoglycemic effect of aqueous extract of the *Anacardium occidentale* Linn leaves grown in Nigeria on normoglycemic albino rats. *JETEAS.* 2012; 3(2): 302-8.
64. Liu RH. Health benefits of fruit and vegetables are from additive and synergistic combinations of phytochemicals. *Am J ClinNutr.* 2003; 78(3): 517S-20S.
65. Brahmabhatt M, Gundala SR, Asif G, Shamsi SA, Aneja R. Ginger phytochemicals exhibit synergy to inhibit prostate cancer cell proliferation. *Nutr Canc.* 2013; 65(2): 263-72.
66. Yao Y, Luong TN, Lepik M, Aftab N, Fong VH, Vieira A. Synergism of antioxidant phytochemicals: comparisons among purified polyphenols and dietary-plant extracts. *ISHS Acta Horticulturae* 939: XXVIII International Horticultural Congress on Science and Horticulture for People (IHCSP) International Symposium on Emerging Health Topics in Fruits and Vegetables; 2010.
67. Kunyanga CN, Imungi JK, Okoth M, Momanyi C, Biesalski HK, Vadivel V. Antioxidant and antidiabetic properties of condensed tannins in acetic extract of selected raw and processed indigenous food ingredients from Kenya. *J Food Sci.* 2011; 76(4): C560-7.
68. Zheng T, Shu G, Yang Z, Mo S, Zhao Y, Mei Z. Antidiabetic effect of total saponins from *Entadaphaseoloides* (L.) Merr in type 2 diabetic rats. *J Ethnopharmacol.* 2012; 139(3): 814-21.
69. Koneri RB, Samaddar S, Ramaiah CT. Antidiabetic activity of a triterpenoidsaponin isolated from *Momordica cymbalaria* Fenzl. *Indian J Exp Biol.* 2014; 52(1): 46-52.
70. Mathew JE, Mantri A, Vachala SD, Srinivasan KK, Movaliya V. Effect of *Sphaeranthus indicus* ethanol extract on tissue antioxidant activity in gentamicin induced nephrotoxic rats. *Herba Polonica.* 2009; 55(4): 86-95.
71. Mordi JC, Awire IE, Awhin PE. Effects of the aqueous extract of *Hibiscus sabdariffa* L petal (zobo) drink on ethanol-induced changes in plasma superoxide dismutase and lipid peroxidation levels in brain and hepatic tissues of rats. *J Biological Sci Bio conserv.* 2011; 3: 35-41.
72. Roopha PD, Padmalatha C. Effect of herbal preparation on heavy metal (Cadmium) induced antioxidant system in female Wistar rats. *J Med Toxicol.* 2012; 8(2): 101–7.
73. Sahreen S, Khan MR, Khan RA, Shah NA. Effect of *Carissa opacaleaves* extract on lipid peroxidation, antioxidant activity and reproductive hormones in male rats. *Lipids Health Dis.* 2013; 12(1): 90. doi: 10.1186/1476-511X-12-90.
74. Sindhu RK, Koo J-R, Roberts CK, Vaziri ND. Dysregulation of hepatic superoxide dismutase, catalase and glutathione peroxidase in diabetes: response to insulin and antioxidant therapies. *Clin Exp Hypertens.* 2004; 26(1): 43–53.

75. Shanmugam KR, Mallikarjuna K, Nishanth K, Kuo CH, Reddy KS. Protective effect of dietary ginger on antioxidant enzymes and oxidative damage in experimental diabetic rat tissues. *Food Chem.* 2011; 124(4): 1436–42.
76. Loven D, Sched H, Wilson H, Daabees TT, Stegink LD, Diekus M, *et al.* Effect of insulin and oral glutathione on glutathione levels and superoxide dismutase activities in organs of rats with streptozocin-induced diabetes. *Diabetes* 1986; 35(5): 503-7.
77. Elmalí E, Altan N, Bukan N. Effect of the sulphonylurea glibenclamide on liver and kidney antioxidant enzymes in streptozocin-induced diabetic rats. *Drugs R & D.* 2004; 5(4): 203-8.
78. Hwang I, Lee J, Huh JY, Park J, Lee HB, Ho Y-S, *et al.* Catalase deficiency accelerates diabetic renal injury through peroxisomal dysfunction. *Diabetes* 2012; 61(3): 728-38.
79. Hünkar T, Aktan F, Ceylan A, Karasu C. Effects of cod liver oil on tissue antioxidant pathways in normal and streptozocin-diabetic rats. *Cell Bio chem Funct.* 2002; 20(4): 297-302.
80. Mieyal JJ, Gallogly MM, Qanungo S, Sabens EA, Shelton MD. Molecular mechanisms and clinical implications of reversible protein S-glutathionylation. *Antiox Redox Signal.* 2008; 10(11): 1941–88.
81. De Mattia G, Bravi MC, Laurenti O, Cassone-Faldetta M, Armiento A, Ferri C, *et al.* Influence of reduced glutathione infusion on glucose metabolism in patients with non-insulin-dependent diabetes mellitus. *Metab.* 1998; 47(8): 993-97.
82. El-Missiry MA, El Gindy AM. Amelioration of alloxan induced diabetes mellitus and oxidative stress in rats by oil of *Eruca sativa* seeds. *Ann Nutri Metabs.* 2000; 44: 97–100.
83. Zitka O, Skalickova S, Gumulec J, Masarik M, Adam V, Hubalek J, *et al.* Redox status expressed as GSH:GSSG ratio as a marker for oxidative stress in paediatric tumor patients. *Oncol Lett.* 2012; 4(6): 1247-53.
84. Murakami K, Takahito K, Ohtsuka Y, Fujiwara Y, Shimada M, Kawakami Y. Impairment of glutathione metabolism in erythrocytes from patients with diabetes mellitus. *Metab.* 1989; 38(8): 753–8.