

Medicinal Plant Images

Ian E Cock^{1,2,*}

¹Centre for Planetary Health and Food Security, Nathan Campus, Griffith University, Nathan, Brisbane, Queensland, AUSTRALIA.

²School of Environment and Science, Nathan Campus, Griffith University, Nathan, Brisbane, Queensland, AUSTRALIA.

Correspondence:

Dr. Ian E Cock^{1,2}

¹Centre for Planetary Health and Food Security, Nathan Campus, Griffith University, 170 Kessels Rd, Nathan, Brisbane, Queensland 4111, AUSTRALIA.

²School of Environment and Science, Nathan Campus, Griffith University, 170 Kessels Rd, Nathan, Brisbane, Queensland 4111, AUSTRALIA.

Email id: i.cock@griffith.edu.au



Figure 1: *Eremophila arachnoides* subsp. *arachnoides* Chinnock (family Scrophulariaceae) is an endemic Australian plant. Several *Eremophila* spp. were used in traditional First Australian medicine to treat a wide variety of ailments.^{1,2} Additionally, many other members of the genus *Eremophila* also have traditional medicinal uses. The genus consists of more than 200 species that grow in semi-arid and arid regions of Australia. Multiple *Eremophila* spp. are used as traditional medicines by the First Australians in the areas in which they grow to treat diverse conditions including uses as antibacterial, antifungal, antiviral, antioxidant, anti-diabetic, and anti-inflammatory therapies, as well as for their cardio-protective properties. The antibacterial properties of *Eremophila* spp. have been relatively well studied and the several bioactive terpenoids have been identified. This photograph was taken in the Australian Arid Lands Botanic Garden, Port Augusta, Australia, in January 2021 by Ian Cock.



Figure 2: *Syzygium australe* (H.L. Wendl. ex-Link) B. Hyland (commonly known as brush cherry). *Syzygium* is a large genus of evergreen flowering plants of the family Myrtaceae which consists of approximately 500 species.³ Plants of this genus are widespread, occurring in tropical and subtropical regions of South-East Asia, Australia and Africa. Many *Syzygium* species produce edible fruits and berries. Some *Syzygium* spp. are used in traditional medicine to treat respiratory ailments, tuberculosis, gastrointestinal disorders, diarrhoea and dysentery.⁴ Recent reports have also highlighted the Australian *Syzygium S. leuhmannii* and *Syzygium australe* (Brush Cherry) extracts as having exceptionally high antioxidant contents.⁵ Antioxidants have been associated with the prevention of cancer, cardiovascular disease and neurological degenerative disorders.⁶⁻⁸ They are also linked with anti-diabetic bioactivities and have been associated with the reduction of obesity. Antioxidants can directly scavenge free radicals, protecting cells against oxidative stress related damage to proteins, lipids and nucleic acids.⁹ Thus, *Syzygium* spp. have potential in the treatment of a significant number of diseases and medical conditions related to cellular redox state.

Many other *Syzygium* species internationally also have documented uses in traditional medicine.³ In the commercially most important species *Syzygium aromaticum* (clove), the unopened flower bud is used as a spice. This plant also has uses in traditional medicine due to its anaesthetic properties.¹⁰ The antibacterial activity of *S. aromaticum* is also well known. Numerous studies have reported on the anti-bacterial¹⁰ and anti-fungal¹¹ activities of oils and extracts from this plant. Other *Syzygium* species from South East Asia (*Syzygium jambos*)¹² and India (*Syzygium lineare* and *Syzygium cumini*)¹³ have also been shown to have anti-microbial activity. Recent studies have also reported the anti-bacterial activity of *Syzygium cordatum* leaf extracts. Of particular interest was the potent growth inhibitory of the extracts against the bacterial triggers of rheumatoid arthritis¹⁴ and ankylosing spondylitis.¹⁵



DOI: 10.5530/pc.2023.4.31

Copyright Information :

Copyright Author (s) 2023 Distributed under Creative Commons CC-BY 4.0

Publishing Partner : EManuscript Tech. [www.emanuscript.in]

REFERENCES

- Cock IE, Baghtchedjian L, Cordon ME, Dumont E. Phytochemistry, medicinal properties, bioactive compounds, and therapeutic potential of the genus *Eremophila* (Scrophulariaceae). *Molecules*. 2022;27(22):7734. doi: 10.3390/molecules27227734, PMID 36431834.
- Cock IE. Medicinal and aromatic plants – Australia. Available from: <http://www.eolss.net>. In: *Ethnopharmacology, Encyclopedia of Life Support Systems (EOLSS)*. Oxford, UK: EOLSS Publishers; 2011. Developed under the auspices of UNESCO.
- Cock IE, Cheesman M. Plants of the genus *Syzygium* (Myrtaceae): a review on ethnobotany, medicinal properties, and phytochemistry. In: *Bioactive compounds of medical plants. Properties and potential for human health*. NJ: Apple Academic Press; 2018.
- Cock IE. Medicinal and aromatic plants – Australia. Available from: <http://www.eolss.net>. In: *Ethnopharmacology, Encyclopedia of Life Support Systems (EOLSS)*, Developed under the auspices of UNESCO. Oxford, UK: EOLSS Publishers.
- Netzel M, Netzel G, Tian Q, Schwartz S, Konczak I. Native Australian fruits – a novel source of antioxidants for food. *Innov Food Sci Emerg Technol*. 2007;8(3):339-46. doi: 10.1016/j.ifset.2007.03.007.
- Potter JD. Cancer prevention: epidemiology and experiment. *Cancer Lett*. 1997;114(1-2):7-9. doi: 10.1016/s0304-3835(97)04615-6, PMID 9103244.
- Cock IE. The medicinal properties and phytochemistry of plants of the genus *Terminalia* (Combretaceae). *Inflammopharmacology*. 2015;23(5):203-29. doi: 10.1007/s10787-015-0246-z, PMID 26226895.
- Cock IE. Problems of reproducibility and efficacy of bioassays using crude extracts, with reference to *Aloe vera*. *Pharmacogn Commun*. 2011;1(1):52-62. doi: 10.5530/pc.2011.1.3.
- Rice-Evans C, Miller N, Paganga G. Antioxidant properties of phenolic compounds. *Trends Plant Sci*. 1997;2(4):152-9. doi: 10.1016/S1360-1385(97)01018-2.
- Arora DS, Kaur GJ. Antibacterial activity of some Indian medicinal plants. *J Nat Med*. 2007;61(3):313-7. doi: 10.1007/s11418-007-0137-8.
- Park MJ, Gwak KS, Yang I, Choi WS, Jo HJ, Chang JW, et al. Antifungal activities of the essential oils of *Syzygium aromaticum* (L.) Merr. Et Perry and *Leptospermum petersonii* Bailey and their constituents against various dermatophytes. *J Microbiol*. 2007;45(5):460-5. PMID 17978807.
- Mohanty S, Cock IE. Bioactivity of *Syzygium jambos* methanolic extracts: antibacterial activity and toxicity. *Pharmacogn Res*. 2010;2(1):4-9. doi: 10.4103/0974-8490.60577, PMID 21808530.
- Duraipandian V, Ayyanar M, Ignacimuthu S. Antimicrobial activity of some ethnomedicinal plants used by Paliyar tribe from Tamil Nadu, India. *BMC Complimentary Altern Med*. 2006;6(35):1-7.
- Cock IE, van Vuuren SF. Anti-Proteus activity of some South African medicinal plants: their potential for the treatment and prevention of rheumatoid arthritis. *Inflammopharmacology*. 2014;22(1):23-36. doi: 10.1007/s10787-013-0179-3, PMID 23877712.
- Cock IE, Van Vuuren SF. The potential of selected South African plants with anti-Klebsiella activity for the treatment and prevention of ankylosing spondylitis. *Inflammopharmacology*. 2015;23(1):21-35. doi: 10.1007/s10787-014-0222-z, PMID 25412961.